



Effective action of string theory at order α in the presence of boundary

Mohammad R. Garousi¹

Department of Physics, Faculty of Science, Ferdowsi University of Mashhad, P.O. Box 1436, Mashhad, Iran

Received: 9 August 2021 / Accepted: 19 December 2021 / Published online: 31 December 2021

© The Author(s) 2021

Abstract Recently, using the assumption that the string theory effective action at the critical dimension is background independent, the classical on-shell effective action of the bosonic string theory at order α in a spacetime manifold without boundary has been reproduced, up to an overall parameter, by imposing the $O(1,1)$ symmetry when the background has a circle. In the presence of the boundary, we consider a background which has boundary and a circle such that the unit normal vector of the boundary is independent of the circle. Then the $O(1,1)$ symmetry can fix the bulk action without using the lowest order equation of motion. Moreover, the above constraints and the constraint from the principle of the least action in the presence of boundary can fix the boundary action, up to five boundary parameters. In the least action principle, we assume that not only the values of the massless fields but also the values of their first derivatives are arbitrary on the boundary. We have also observed that the cosmological reduction of the leading order action in the presence of the Hawking–Gibbons boundary term, produces zero cosmological boundary action. Imposing this as another constraint on the boundary couplings at order α , we find the boundary action up to two parameters. For a specific value for these two parameters, the gravity couplings in the boundary become the Chern–

Simons gravity plus another term which has the Laplacian of the extrinsic curvature.

1 Introduction

String theory is a quantum theory of gravity with a finite number of massless fields and a tower of infinite number of massive fields reflecting the stringy nature of the gravity. The critical dimension for the bosonic string is 26, and for the type IIA, type IIB, type I and for the heterotic strings is 10. The type IIB superstring theory on a spacetime manifold with negative cosmological constant which has boundary is conjectured to be dual to a conformal field theory on the boundary [1,2]. The string theory is usually explored by studying its effective action which includes the massless fields and their higher derivative terms. For the spacetime manifolds with boundary, the effective action has both bulk and boundary terms, i.e., $S_{\text{eff}} + \partial S_{\text{eff}}$. At the leading order of the derivative, the bulk action should include the Hilbert–Einstein action at the critical dimension and the boundary action should include the corresponding Hawking–Gibbons–York boundary term [3,4]. These actions and their appropriate higher derivative extensions should be produced by specific techniques in the string theory.

The effective actions in the string theory have a double expansions. The genus-expansion which includes the classical tree-level and a tower of quantum loop-level corrections, and the stringy-expansion which is an

¹ e-mail: garousi@um.ac.ir (corresponding author)

expansion in terms of higher derivative couplings at each loop level. It has been shown in [5–8] that the tree-level effective action of the bosonic string theory at orders α^0 and α are invariant under $O(d,d)$ transformations if one compactifies the theory on the torus T^d and keeps only the zero modes (cosmological reduction). Using the string field theory, it has been proved in [9] that the cosmological reduction of the tree-level effective action of the bosonic string theory to all orders of α should be invariant under $O(d,d)$ transformations. This has been extended in [10] to the classical effective action of the heterotic string theory.

The Einstein theory of general relativity is background independent in the sense that only gauge symmetry is required to specify the theory. We expect that the string theory classical effective action at the critical dimension which is a higher-derivative extension of the Einstein theory at the critical dimension, to be background independent too. Unlike the Einstein action which has only one coupling, however, there are many gauge invariant couplings in the effective action of the string theory at each order of α , e.g., at the leading order the independent gauge invariant couplings in the bosonic string theory are

$$S_0 = -\frac{2}{\kappa^2} \int d^{26}x e^{-2\Phi} \sqrt{-G} \times (\alpha_1 R + \alpha_2 \nabla_a \Phi \nabla^a \Phi + \alpha_3 H^2), \quad (1)$$

where $\alpha_1, \alpha_2, \alpha_3$ are three parameters. The first term is the Einstein action at the critical dimension 26. The background independence assumption then requires these parameters to be independent of the geometry of the spacetime, i.e., if the background has the torus T^4 or K^3 the value of the coefficients $\alpha_1, \alpha_2, \alpha_3$ remains the same. However, the coefficients of the gauge invariant couplings in the reduced action do depend on the geometry of the compact spaces. In other words, if one compactifies the above action on T^4 , the result would be the same as the compactification on K^3 or any other

compact manifold provided that one takes into account all the corresponding Kaluza–Klein modes. However, if one ignores the Kaluza–Klein modes (dimensional reduction), then the actions in the lower dimension have different symmetries corresponding to the compact spaces, i.e., the 22-dimensional action in the case of T^4 has symmetry $O(4,4)$ which is different than the symmetry of the 22-dimensional action in the case of K^3 . This means, if one could fix somehow the parameters of the effective action at the critical dimension for a particular geometry in which the reduced action has a specific symmetry, then that parameters would be valid for any other geometry. For example, if one considers the

background to have a circle, then the dimensional reduction of the action should have the $O(1,1)$ symmetry. This symmetry has been used in [11] to fix the parameters in the above action up to an overall factor, i.e.,

$$S_0 = -\frac{2\alpha_1}{\kappa^2} \int d^{26}x e^{-2\Phi} \sqrt{-G} \times \left(R + 4\nabla_a \Phi \nabla^a \Phi - \frac{1}{12} H^2 \right), \quad (2)$$

which is the standard effective action of the bosonic string theory for $\alpha_1 = 1$. At the higher orders of α , there is the complication that the effective action has the freedom of the higher-derivative field redefinitions [12]. In these cases, the $O(1,1)$ symmetry may fix the parameters of the independent gauge invariant couplings up to the field redefinitions

When the geometry has one circle, the constraints from the Z_2 -subgroup of the $O(1,1)$ symmetry have been used in [11,13] to find the effective actions of the bosonic string theory at four- and six-derivative orders in a minimal scheme, up to an overall factor. Assuming there is such symmetry for the classical effective action of the type II superstring theories as well, all eight-derivative couplings for NS–NS fields have been found in [14–17], up to an overall factor. The background independent assumption then requires the resulting couplings to be valid for any other spacetime, up to the field redefinitions. In fact, the effective actions found in this way are fully consistent with the sphere-level S-matrix element of four NS–NS vertex operators and with the results from the sigma model [16,17]. Moreover, when the geometry has the torus T^d , the cosmological reduction of these effective actions are also fully consistent with the $O(d,d)$ symmetry [18,19]. Assuming also the classical world-volume effective actions of the non-perturbative branes in the string theory transform covariantly under the Z_2 -transformations when the spacetime geometry has a circle, then many already known and unknown world-volume couplings have been found in [20,21].

In applying the $O(1,1)$ symmetry to find the couplings at orders $\alpha, \alpha^2, \alpha^3$ in [11,13–17], one first needs to find all *independent* gauge invariant couplings in the minimal scheme, i.e., the couplings which are not related by various Bianchi identities, by the field redefinitions and by total derivative terms. The number of independent couplings at each order of α is fixed, however, the structure of the gauge invariant couplings depends on how to use the above freedom to find the independent couplings. The number of independent couplings at orders $\alpha^1, \alpha^2, \alpha^3$ in the bosonic theory are 8, 60, 872, respectively. These couplings in a specific minimal

scheme have been found in [12,15,22]. Then one should impose the Z_2 -symmetry on these independent couplings to find their corresponding coefficients at each order of α up to one parameter. The Z_2 -transformations or T-duality transformations are the Buscher rules [23,24] and some higher derivative corrections at each order of α which depends on the minimal scheme that one uses for the gauge invariant couplings at that order [11]. If one does not use the field redefinitions to write the gauge invariant couplings in the minimal scheme, then the $O(1,1)$ constraint can fix the effective action up to many parameters which can be removed by the field redefinition [11]. These parameters appear also in the corrections to the Buscher rules. By changing these corrections, one can change the scheme of the gauge invariant couplings. However, there would be no scheme for which the T-duality transformations are only the Buscher rules.

When the spacetime geometry has one circle, the Z_2 symmetry imposes the following constraint on the bulk effective action:

$$S_{\text{eff}}(\psi) = S_{\text{eff}}(\psi') \tag{3}$$

where S_{eff} is the reduction of S_{eff} , ψ represents the massless fields in the base space and ψ is its Z_2 -transformations. There are always some total derivative terms in the base space [14–17] which become zero when spacetime has no boundary. However, when the spacetime has boundary, the presence of the total derivative terms dictates that there must be some couplings on the boundary as well.

For the spacetime manifold which has boundary, using the background independent assumption, one may consider a geometry that has a boundary and one circle. Then the Z_2 symmetry may fix the couplings in the bulk and boundary actions up to field redefinitions. It has been speculated in [25] that, in the presence of the boundary, the invariance of the classical effective action under the Z_2 -transformations should be extended as follows: The sum of the bulk and the boundary actions, i.e., $S_{\text{eff}} + \partial S_{\text{eff}}$, should be invariant under the Z_2 -transformations, i.e.,

$$S_{\text{eff}}(\psi) + \partial S_{\text{eff}}(\psi) = S_{\text{eff}}(\psi') + \partial S_{\text{eff}}(\psi') \tag{4}$$

where ∂S_{eff} is the reduction of the boundary action ∂S_{eff} . There might be some total derivative terms on the boundary of the base space, however, they become zero using the Stokes's theorem because the boundary of boundary is zero. In this paper, we are going to impose the above constraint on the effective action of the bosonic string theory at order α . We consider the background that its boundary is independent of the circle, i.e., the unit normal vector to the boundary is invariant under the T-duality transformations at order α . This particular background constrains the corrections to

the Buscher rules. We will see that for this Z_2 -transformations, the $O(1,1)$ constraint is not consistent with the effective action in the minimal scheme. In fact to impose the $O(1,1)$ symmetry for this geometry one should not use the field redefinitions to reduce the number of gauge invariant couplings at order α .

The constraint (4) has been used in [26] to find the O-plane effective action at order α^2 in the presence of the boundary in the type II superstring theories in which the Buscher rules have no correction at orders α^1, α^2 . The constraint (4) has been also used in [25] to find the spacetime effective action of the bosonic string theory at order α^1 . This constraint fixes the bulk actions completely, however, it fixes the boundary action up to some extra boundary parameters.

When spacetime has boundary, however, there are further constraints on the boundary actions from the principle of the least action. To be able to extremize the bulk effective actions at each order of α , the boundary should have specific couplings and the massless fields should have appropriate values on the boundary. For example, the Hawking–Gibbons boundary term along with the arbitrariness of the metric on the boundary is needed to be able to extremize the Hilbert–Einstein action, i.e., the Einstein's equations are derived by extremizing the Hilbert–Einstein action against variations of the spacetime metric $G_{\mu\nu}$ which is arbitrary on the boundary, i.e., $\delta G_{\mu\nu}$ and its tangent derivatives along the boundary are zero. The normal derivative of the metric, however, is not arbitrary on the boundary. The variation of this term which is not zero, appears on the boundary when one extremizes the bulk action. The variation of the Hawking–Gibbons boundary term cancels this normal derivative term on the boundary. We expect similar constraint for the boundary terms in the effective actions of the string theory. However, if one assumes only the metric is arbitrary on the boundary, the standard gravity couplings in the effective actions of the string theory at orders α^2, α^3 cannot be extremized for any boundary couplings. In fact it has been shown in [27] that only the gravity couplings in the Euler character can be extremized. We will show that the $O(1,1)$ -constraint on the couplings at order α produces the bulk gravity couplings which are the same as the bulk couplings in the Euler character, however, it produces the boundary couplings which are consistent with the Chern–Simons form as well as some other gravity coupling in the boundary.

Hence, in order to be able to extremize the effective action of the string theory at order α^n , we propose that not only the massless fields but also their derivatives up to order n should be arbitrary on the boundary, i.e., the massless

fields are arbitrary on the boundary for the effective action at order α^0 , the massless fields and their first derivatives are arbitrary on the boundary for the effective action at order α , and so on. This may be inspired by the fact that the linear differential equation $d^{2n+2} x_n(t) = 0$ has specific solution when the functions $x, \frac{dx}{dt}, \dots, \frac{d^{2n}x}{dt^{2n}}$ are known at the initial and the final times.

Imposing the $O(1,1)$ -symmetry on the most general gauge invariant couplings at order α^0 , one finds that the effective actions of the bosonic string theory are fixed up to one extra parameter in the boundary action [25], i.e.,

$$\begin{aligned} \mathbf{S}_0 + \partial \mathbf{S}_0 = & -\frac{2\alpha_1}{\kappa^2} \left[\int d^D x \sqrt{-G} e^{-2\Phi} \right. \\ & \times \left(R + 4\nabla_\mu \Phi \nabla^\mu \Phi - \frac{1}{12} H^2 \right) \\ & \left. + 2 \int d^{D-1} \sigma \sqrt{|g|} e^{-2\Phi} K \right] \\ & - \frac{2\alpha_5}{\kappa^2} \int d^{D-1} \sigma e^{-2\Phi} \sqrt{|g|} \left(-\frac{1}{2} K + n^\mu \nabla_\mu \Phi \right) \end{aligned} \quad (5)$$

where K is trace of the extrinsic curvature and n^μ is normal vector to the boundary. It is outward-pointing (inward-pointing) if the boundary is spacelike (timelike). We consider in this paper only the timelike boundary. The above actions are invariant under the Z_2 -transformation for arbitrary parameters α_1, α_5 . The sum of the bulk and boundary terms in the first line are Z_2 -invariant, and the boundary terms in the second line are also invariant under the Z_2 transformations. The standard normalization of the Einstein term fixes $\alpha_1 = 1$. However, the parameters α_5 remains arbitrary. In the superstring theory, there are S-duality as well which constrains the parameter α_5 to be zero [25]. In the bosonic string theory, however, there is no such symmetry. One can fix this parameter by the principle of the least action as follows: Since the action is at two derivative order, only the massless fields are arbitrary on the boundary. In extremizing the bulk action, the normal derivative of the variation of dilaton which is not zero, does not appear on the boundary, whereas, the variation of the boundary action in the second line above produces such a term. The only way to cancel this term, i.e., to be able to extremize the bulk and boundary actions, is to set $\alpha_5 = 0$. Hence, the constraints from the Z_2 -symmetry and the least action principle, reproduce the standard bulk and boundary actions at the leading order of α , i.e.,

$$\mathbf{S}_0 + \partial \mathbf{S}_0 = -\frac{2}{\kappa^2} \left[\int d^D x \sqrt{-G} e^{-2\Phi} \right.$$

$$\begin{aligned} & \left. \times \left(R + 4\nabla_\mu \Phi \nabla^\mu \Phi - \frac{1}{12} H^2 \right) \right. \\ & \left. + 2 \int d^{D-1} \sigma \sqrt{|g|} e^{-2\Phi} K \right] \end{aligned} \quad (6)$$

However, it turns out that if one imposes the Z_2 -constraint (4) and the constraint from the least action principle to the effective actions of the bosonic string theory at order α , one can not fully fix all parameters in the boundary action.

The cosmological reduction of the classical bulk actions must be invariant under $O(d,d)$ transformations [9]. We expect the cosmological reduction of the boundary actions should be also invariant under the $O(d,d)$ transformations. In the observation that the cosmological reduction of the leading order bulk action is invariant under the $O(d,d)$ symmetry, one removes a total derivative term which is not invariant under the $O(d,d)$ transformations. We find that the cosmological reduction of the Hawking–Gibbons term is not invariant under the $O(d,d)$ transformations either. However, if one keeps track of the total derivative term and transfers it to the boundary by using the Stokes's theorem, one observes that the cosmological reduction of the boundary action at the leading order becomes invariant under the $O(d,d)$ transformations. In fact it becomes zero. This motivates us to speculate that the cosmological reduction of the classical boundary actions at any order of α must be invariant under the $O(d,d)$ transformations, and may even be zero, i.e.,

$$\partial \mathbf{S}_{\text{eff}}^\alpha = 0 \quad (7)$$

In this paper, we would like to impose the Z_2 -constraint (4), the constraint from the least action principle and the above constraint on the cosmological reduction of the boundary actions, to fix the effective actions of the bosonic string theory at order α when the spacetime has boundary.

The outline of the paper is as follows: in Sect. 2, we use the Bianchi identities and remove the total derivative terms from the bulk action to the boundary action to show that there are 20 independent bulk and 38 independent boundary gauge invariant couplings at order α , without using the field redefinitions. In Sect. 3, using the background independent assumption, we consider a specific background geometry which has a circle and a boundary that its normal vector is independent of the circle. Then using the fact that the circle reduction of the effective action on this background should have $O(1,1)$ symmetry, we constrain the coefficients of the couplings. In Sect. 3.1, we show that the T-duality constraint in the bulk fixes the 20 parameters in terms of two parameters. We impose a relation between these two parameters by requiring the effective action to have the standard propagator for the B -field. The resulting bulk action

is exactly the one found by K.A. Meissner up to one overall factor. The Tduality constraint on the bulk couplings produces also some total derivative terms in the base space which are transferred to the boundary by using the Stokes’s theorem. In Sect. 3.2, we show that the T-duality constraint on the boundary couplings fixes the 38 boundary parameters in terms of the overall bulk factor and in terms of 7 boundary parameters. In Sect. 4, we impose the constraint from the principle of the least action. Since the independent bulk couplings have no term with three derivatives, extremizing the bulk action produces no constraint on the bulk parameters. However, extremizing the boundary action, one can fix 2 of the 7 boundary parameters. In Sect. 5, we study the Cosmological/one-dimensional reduction of the actions. We find that the cosmological constraint (7) on the boundary action fixes 3 of the 5 boundary parameters. In Sect. 6, we briefly discuss our results.

2 Gauge invariance constraint at order α

The effective action of the string theory has a double expansions. One expansion is the genus expansion which includes the classical sphere-level and a tower of quantum effects. The other one is the stringy expansion which is an expansion in terms of higher-derivative couplings. The number of derivatives in each coupling can be accounted by the order of α . When spacetime has boundary, the sphere-level effective action $S_{\text{eff}} + \partial S_{\text{eff}}$ has the following α -expansion in the string frame:

$$S_{\text{eff}} = \sum_m \alpha^m S_m = S_0 + \alpha' S_1 + \dots \quad (8)$$

$$\partial S_{\text{eff}} = \sum_m \alpha^m \partial S_m = \partial S_0 + \alpha' \partial S_1 + \dots ;$$

where G is determinant of the bulk metric $G_{\mu\nu}$ and boundary is specified by the functions $x^\mu = x^\mu(\sigma^\mu)$. In the second line, g is determinant of the induced metric on the boundary

$$g = \det \left(\frac{\partial x^\mu}{\partial \sigma^{\tilde{\mu}}} \frac{\partial x^\nu}{\partial \sigma^{\tilde{\nu}}} G_{\mu\nu} \right) \partial \mu \partial \nu$$

$$g_{\mu\nu} = \dots \quad (9)$$

The effective action must be invariant under the coordinate transformations and under the B -field gauge transformations. One can easily find the independent couplings in the bulk and boundary actions at order α^0 , i.e.,

$$\begin{aligned} \mathcal{L}_0 &= \alpha_1 R + \alpha_2 \nabla_\mu \Phi \nabla^\mu \Phi + \alpha_3 H^2 \\ \partial \mathcal{L}_0 &= \alpha_4 K + \alpha_5 n^\mu \nabla_\mu \Phi \end{aligned} \quad (10)$$

where $\alpha_1, \dots, \alpha_5$ are 5 parameters that the gauge symmetry cannot fix them. Using the background independent assumption, they can be fixed by the Z_2 -symmetry (5), and by the least action principle (6).

Using the package “xAct” [28], one finds there are 41 gauge invariant couplings in the bulk action at order α . However, they are not all independent. To find the independent bulk couplings, we note that the total derivative terms in the bulk can be transferred to the boundary using the Stoke’s theorem. Hence, the couplings in the bulk should not include total derivative terms. Moreover, the independent couplings should not be related to each others by the Bianchi identities

$$\begin{aligned} R_{\alpha[\beta\gamma\delta]} &= 0 \\ \nabla_{[\mu} R_{\alpha\beta]\gamma\delta} &= 0 \\ \nabla_{[\mu} H_{\alpha\beta\gamma]} &= 0 \\ [\nabla, \nabla]0 &- R0 = 0 \end{aligned} \quad (11)$$

Removing the above freedoms from the most general gauge invariant couplings in the bulk action, one finds there are 20 even-parity independent couplings [12], i.e.,

$$\begin{aligned} \mathcal{L}_1 &= a_7 R^2 + a_6 R H_{\alpha\beta\gamma} H^{\alpha\beta\gamma} + a_1 H_\alpha^{\delta\epsilon} H^{\alpha\beta\gamma} H_{\beta\delta}^\epsilon H_{\gamma\epsilon\epsilon} \\ &\quad + a_2 H_{\alpha\beta\gamma} H_{\gamma}^{\epsilon\epsilon} H_{\delta\epsilon\epsilon} + a_2 H_{\alpha\beta} \\ &\quad + a_3 H_{\alpha\beta\gamma} H_{\alpha\beta\gamma} H^{\delta\epsilon\epsilon} H^{\delta\epsilon\epsilon} + a_4 H_\alpha^{\gamma\delta} H_{\beta\gamma\delta} R^{\alpha\beta} \\ &\quad + a_5 R_{\alpha\beta} R_{\alpha\beta} + a_8 R_{\alpha\beta\gamma\delta} R_{\alpha\beta\gamma\delta} \\ &\quad + a_9 H_\alpha^{\delta\epsilon} H_{\alpha\beta\gamma} H_{\beta\gamma\delta} + a_{11} R \nabla_\alpha \nabla^\alpha \Phi \\ &\quad + a_{10} H_{\beta\gamma\delta} H^{\beta\gamma\delta} \nabla_\alpha \nabla^\alpha \Phi + a_{13} R \nabla_\alpha \Phi \nabla^\alpha \Phi \\ &\quad + a_{12} H_{\beta\gamma\delta} H^{\beta\gamma\delta} \nabla_\alpha \Phi \nabla^\alpha \Phi + a_{14} \nabla_\alpha \Phi \nabla^\alpha \Phi \nabla_\beta \nabla^\beta \Phi \\ &\quad + a_{15} H_\alpha^{\gamma\delta} H_{\beta\gamma\delta} \nabla^\alpha \Phi \nabla^\beta \Phi \\ &\quad + a_{16} R_{\alpha\beta} \nabla^\alpha \Phi \nabla^\beta \Phi + a_{17} \nabla_\alpha \Phi \nabla^\alpha \Phi \nabla_\beta \Phi \nabla^\beta \Phi \\ &\quad + a_{18} H_\alpha^{\gamma\delta} H_{\beta\gamma\delta} \nabla^\beta \nabla^\alpha \Phi \\ &\quad + a_{19} \nabla_\beta \nabla_\alpha \Phi \nabla^\beta \nabla^\alpha \Phi \\ &\quad + a_{20} \nabla_\alpha H^{\alpha\beta\gamma} \nabla_\delta H_{\beta\gamma}^\delta \end{aligned} \quad (12)$$

where a_1, \dots, a_{20} are 20 parameters that the gauge symmetry can not fix them. The assumption that the effective action is

background independent means these parameters are background independent. They may be fixed for the particular

geometry which has one circle. Note that the above couplings have no term with three derivatives, hence, in extremizing the above Lagrangian one does not face with the variation of the second derivative of massless fields on the boundary which are non-zero. As a result, our proposal for the boundary conditions in which the massless fields and their first derivatives are arbitrary on the boundary, i.e., their variations are zero on the boundary, does not constraint the parameters a_1, \dots, a_{20} . In other words, the above bulk action satisfies $\delta S_1 = 0$ for any values of the parameters.

Since the boundary of spacetime has a unit normal vector n^μ , the boundary Lagrangian ∂L_1 should include this vector as well as the tensors $K_{\mu\nu}$, $H_{\mu\nu\rho}$, $R_{\mu\nu\rho\sigma}$, $\nabla_\mu \Phi$ and their derivatives at order α . The second fundamental form or the extrinsic curvature of boundary, i.e., $K_{\mu\nu}$, is defined as $K_{\mu\nu} = P_{\mu}^{\alpha} P_{\nu}^{\beta} \nabla_{(\alpha} n_{\beta)}$ where $P^{\mu\nu} = G^{\mu\nu} - n^\mu n^\nu$ is the first fundamental form which projects the spacetime tensors tangent to the boundary. Using the fact that n^μ is unit vector orthogonal to the boundary, one can write it as

$$n^\mu = (\partial_\alpha f \partial^\alpha f)^{-1/2} \partial^\mu f \quad (13)$$

where the boundary is specified by the function f to be a constant f^* . One can rewrite $K_{\mu\nu}$ as

$$K_{\mu\nu} = \nabla_{\mu} n_{\nu} - n_{\mu} a_{\nu} \quad (14)$$

where $a_\nu = n^\rho \nabla_{\rho} n_\nu$ is acceleration. It satisfies the relation $n^\mu a_\mu = 0$. Note that the extrinsic curvature is symmetric and satisfies $n^\mu K_{\mu\nu} = 0$ and $n^\mu n^\nu \nabla_{\alpha} K_{\mu\nu} = 0$ which can easily be seen by writing them in terms of function f . Using these symmetries and $n^\mu n_\mu = -1$, one finds there are 56 gauge invariant even-parity couplings in the boundary action, i.e.,

$$\begin{aligned} \partial L_1 = & b'_1 H_{\beta\gamma\delta} H^{\beta\gamma\delta} K^\alpha{}_\alpha + b'_2 H_{\alpha}{}^{\gamma\delta} H_{\beta\gamma\delta} K^{\alpha\beta} \\ & + b'_3 K_{\alpha\gamma} K_{\alpha\beta} K_{\beta\gamma} + b'_4 K_{\alpha\alpha} K_{\beta\gamma} K_{\beta\gamma} \\ & + b'_5 K^\alpha{}_\alpha K^\beta{}_\beta K^\gamma{}_\gamma + b'_6 H_{\alpha}{}^{\delta\epsilon} H_{\beta\delta\epsilon} K^\gamma{}_\gamma n^\alpha n^\beta \\ & + b'_7 H_{\alpha\gamma}{}^\epsilon H_{\beta\delta\epsilon} K_{\gamma\delta} n_{\alpha\beta} + b'_8 K_{\alpha\beta} R_{\alpha\beta\gamma} n_{\alpha\beta} R \\ & + b'_9 K^\gamma{}_\gamma + b K_{\alpha\alpha} R_{\alpha\beta}{}^{10} \\ & + b'_{11} K_{\gamma\delta} n_{\alpha\beta} R_{\alpha\beta\gamma\delta} + b'_{12} H^{\beta\gamma\delta} n^\alpha \nabla_{\alpha} H_{\beta\gamma\delta} \\ & + b'_{13} K_{\beta\gamma} n_{\alpha} \nabla_{\alpha} K_{\beta\gamma} + b'_{14} K^\beta{}_\beta n^\alpha \nabla_{\alpha} K^\gamma{}_\gamma \\ & + b'_{15} n^\alpha \nabla_{\alpha} R + b'_{16} H_{\beta\gamma\delta} H^{\beta\gamma\delta} n^\alpha \nabla_{\alpha} \Phi \\ & + b'_{17} K_{\beta\gamma} K_{\beta\gamma} n_{\alpha} \nabla_{\alpha} \Phi + b'_{18} K^\beta{}_\beta K^\gamma{}_\gamma n^\alpha \nabla_{\alpha} \Phi \end{aligned}$$

$$\begin{aligned} & + b'_{19} H_{\beta}{}^{\delta\epsilon} H_{\gamma\delta\epsilon} \delta_{\alpha\beta} n_{\alpha} n_{\beta} n^\gamma \nabla_{\alpha} \Phi \\ & + b'_{21} n_{\alpha} n_{\beta} n_{\gamma} R_{\beta\gamma}{}^{\nu\alpha} \nabla_{\alpha} \Phi + b'_{22} n^\alpha R \nabla_{\alpha} \Phi \\ & + b'_{23} n^\alpha \nabla_{\alpha} \nabla_{\beta} \nabla^{\beta} \Phi + b'_{24} K^\beta{}_\beta \nabla_{\alpha} \Phi \nabla^{\alpha} \Phi + b'_{25} \nabla_{\alpha} \Phi \nabla_{\beta} K_{\alpha}{}^{\beta} \\ & + b'_{26} n^\alpha n^\beta \nabla_{\alpha} \Phi \nabla_{\beta} K^\gamma{}_\gamma + b'_{27} n^\alpha \nabla_{\beta} R_{\alpha}{}^{\beta} \\ & + b'_{28} K^\gamma{}_\gamma n^\alpha n^\beta \nabla_{\alpha} \Phi \nabla_{\beta} \Phi + b'_{29} \nabla_{\beta} \nabla_{\alpha} K^{\alpha\beta} + b'_{30} n^\alpha n^\beta \nabla_{\beta} \nabla_{\alpha} K^\gamma{}_\gamma + b'_{31} K^{\alpha\beta} \nabla_{\beta} \nabla_{\alpha} \Phi + b'_{32} K^\gamma{}_\gamma n^\alpha n^\beta \nabla_{\beta} \nabla_{\alpha} \Phi + b'_{33} \nabla_{\beta} \nabla^{\beta} K^{\alpha}{}_{\alpha} \\ & + b'_{34} K^{\alpha}{}_{\alpha} \nabla_{\beta} \nabla^{\beta} \Phi + b'_{35} n^\alpha \nabla_{\alpha} \Phi \nabla_{\beta} \nabla^{\beta} \Phi \\ & + b'_{36} n^\alpha \nabla_{\beta} \nabla^{\beta} \nabla_{\alpha} \Phi + b'_{37} n_{\alpha} n_{\beta} \nabla_{\beta} \nabla_{\gamma} K_{\alpha\gamma} \\ & + b'_{38} H_{\alpha}{}^{\gamma\delta} H_{\beta\gamma\delta} n^\alpha \nabla^{\beta} \Phi + b'_{39} n^\alpha R_{\alpha\beta} \nabla^{\beta} \Phi \\ & + b'_{40} K_{\alpha\beta} \nabla^{\alpha} \Phi \nabla^{\beta} \Phi + b'_{41} n^\alpha \nabla_{\alpha} \Phi \nabla_{\beta} \Phi \nabla^{\beta} \Phi \\ & + b'_{42} n^\alpha \nabla_{\beta} \nabla_{\alpha} \Phi \nabla^{\beta} \Phi \\ & + b'_{43} H_{\alpha\delta\epsilon}{}^{\alpha}{}_{\beta} n_{\gamma} n^{\gamma} \nabla_{\gamma} H_{\beta\delta\epsilon} + b'_{44} K^{\beta\gamma} n^{\alpha} \nabla_{\gamma} K_{\alpha\beta} \\ & + b'_{45} K^{\beta}{}_{\beta} n^{\alpha} \nabla_{\gamma} K_{\alpha}{}^{\gamma} \\ & + b'_{46} n_{\alpha} n_{\beta} \nabla_{\alpha} \Phi \nabla_{\gamma} K_{\beta}{}^{\gamma} + b'_{47} n^{\alpha} n^{\beta} n^{\gamma} \nabla_{\gamma} R_{\alpha\beta} \\ & + b'_{48} n_{\alpha} n_{\beta} n^{\gamma} \nabla_{\alpha} \Phi \nabla_{\beta} \Phi \nabla_{\gamma} \Phi \\ & + b'_{49} n_{\alpha} n_{\beta} \nabla_{\gamma} \nabla_{\beta} K_{\alpha\gamma} \\ & + b'_{50} n_{\alpha} n_{\beta} n_{\gamma} \nabla_{\alpha} \Phi \nabla_{\gamma} \nabla_{\beta} \Phi \\ & + b'_{51} n_{\alpha} n_{\beta} n_{\gamma} \nabla_{\gamma} \nabla_{\beta} \nabla_{\alpha} \Phi \\ & + b'_{52} n_{\alpha} n_{\beta} \nabla_{\gamma} \nabla_{\gamma} K_{\alpha\beta} + b'_{53} n^{\alpha} n^{\beta} \nabla_{\beta} K_{\alpha\gamma} \nabla^{\gamma} \Phi \\ & + b'_{54} H^{\beta\gamma\delta} n^{\alpha} \nabla_{\delta}{}^{\alpha\beta\gamma} H \\ & + b'_{55} H_{\alpha}{}^{\beta\gamma} n^{\alpha} \nabla_{\delta} H_{\beta\gamma}{}^{\delta} \\ & + b'_{56} n^{\alpha} n^{\beta} n^{\gamma} n^{\delta} \nabla_{\delta} \nabla_{\gamma} K_{\alpha\beta} \end{aligned} \quad (15)$$

where b_1, \dots, b_{56} are 56 parameters. The terms in the boundary action which have bulk fields and have one vector n^μ can be interpreted as the total derivative terms in the bulk action that are transferred to the boundary by the Stoke's theorem. Note that we have not considered the curvature tensors and the covariant derivatives that are made of the induced metric (9), because they are related to the curvature tensors and covariant derivatives constructed from the spacetime metric by various Gauss–Codazzi relations.

Not all of the couplings in (15) however are independent. Some of them are related by the Bianchi identities and some others by the total derivative terms in the boundary. To remove the

$$\int_{\partial M^{(D)}} \equiv \int_{\partial M^{(D)}} d^{D-1} \sigma \sqrt{|g|} T_1 \quad (16)$$

redundancy corresponding to the total derivative terms, we add to ∂L_1 all total derivative terms at order α with arbitrary coefficients. The total derivative terms in the boundary have different structure than the total derivative terms in the bulk. According to the Stokes's theorem, the total derivative terms in the boundary which have the following structure:

are zero because the boundary of boundary is zero (see e.g., the appendix in [26]). In above equation $\mathcal{F}_1^{\alpha\beta}$ is an arbitrary antisymmetric even-parity tensor constructed from $n, K, \nabla K, H^2, \nabla\Phi, \nabla\nabla\Phi, R$ at two-derivative order, i.e.,

$$\begin{aligned} \mathcal{F}_1^{\alpha\beta} = & f_2(H^{\beta\delta\epsilon} H_{\gamma\delta\epsilon} n^\alpha n^\gamma - H^{\alpha\delta\epsilon} H_{\gamma\delta\epsilon} n^\beta n^\gamma) + f_3(n \\ & + f_4(n^\gamma \nabla^\alpha K^\beta_\gamma - n^\gamma \nabla^\beta K^\alpha_\gamma) n R_\gamma \\ & + f_5(n^\delta \nabla^\alpha K^\beta_\delta - n^\delta \nabla^\beta K^\alpha_\delta) - n n \\ & + f_6(n^\beta \nabla^\alpha K^\gamma_\gamma - n^\alpha \nabla^\beta K^\gamma_\gamma) R_\gamma \\ & + f_7(n^\beta \nabla^\alpha K^\delta_\delta - n^\alpha \nabla^\beta K^\delta_\delta) \\ & + f_8(K^\gamma_\gamma n^\beta \nabla^\alpha \Phi - K^\gamma_\gamma n^\alpha \nabla^\beta \Phi) \\ & + f_9(K^\delta_\delta n^\beta \nabla^\alpha \Phi - K^\delta_\delta n^\alpha \nabla^\beta \Phi) \\ & + f_{10}(n^\beta \nabla_\gamma K^{\alpha\gamma} - n^\alpha \nabla_\gamma K^{\beta\gamma}) \\ & + f_{11}(n^\beta n^\gamma \nabla^\alpha \Phi \nabla_\gamma \Phi - n^\alpha n^\gamma \nabla^\beta \Phi \nabla_\gamma \Phi) \\ & + f_{12}(n^\beta n^\gamma \nabla_\gamma \nabla^\alpha \Phi - n^\alpha n^\gamma \nabla_\gamma \nabla^\beta \Phi) \\ & f_{13}(K^\beta_\gamma n^\alpha \nabla^\gamma \Phi - K^\alpha_\gamma n^\beta \nabla^\gamma \Phi) + \\ & + f_{14}(n\beta n\gamma n\delta \nabla\delta K\alpha\gamma - n\alpha n\gamma n\delta \nabla\delta K\beta\gamma) \\ & + f_{15}(n\beta \nabla\delta K\alpha\delta - n\alpha \nabla\delta K\beta\delta) \\ & + f_{16}(K^\beta_\delta n^\alpha \nabla^\delta \Phi - K^\alpha_\delta n^\beta \nabla^\delta \Phi) \end{aligned} \quad (17)$$

where f_2, \dots, f_{16} are 15 arbitrary parameters. Adding the above total derivative terms to ∂L_1 , one finds the same Lagrangian but with different parameters b_1, b_2, \dots . We call the new Lagrangian ∂L_1 . Hence

$$\Delta T_1 = 0 \quad (18)$$

where $\Delta = \partial \mathcal{L}_1 - \partial L_1$ is the same as ∂L_1 but with coefficients $\delta b_1, \delta b_2, \dots$ where $\delta b_i = b_i - b_i$. Solving the above equation, after imposing the Bianchi identities (11) and the identities corresponding to the unit vector n^μ , one finds some linear relations between only $\delta b_1, \delta b_2, \dots$ which indicate how the couplings are related among themselves by the total derivative terms, by the Bianchi identities, and by the identities corresponding to the unit vector. The above equation also gives some relations between the coefficients of the total derivative terms and $\delta b_1, \delta b_2, \dots$ in which we are not interested.

To impose the Bianchi identities (11) and the identities corresponding to the unit vector n , we write the covariant derivatives and the curvatures in terms of partial derivatives of metric, dilaton, H and n . We then write the partial derivatives of H in terms of potential B -field and the partial derivatives of n in terms of function f , using the definition (13). Then all the Bianchi identities (11) and the identities corresponding to the unit vector n satisfy automatically. In other words, there is no identities any more when one rewrites everything in terms of metric, dilaton, the potential B -field and f . To simplify the calculation, one may go to the local frame [22] in which the first partial derivative of metric is zero. All these steps can be done easily by the computer. Then one finds 38 relations between only $\delta b_1, \delta b_2, \dots$ which indicate there are 38 independent couplings. One particular choice for the independent boundary couplings is the following:

$$\begin{aligned} \partial L_1 = & b_1 H_{\beta\gamma\delta} H_{\beta\gamma\delta} K_{\alpha\alpha} + b_2 H_{\alpha\gamma\delta} H_{\beta\gamma\delta} K_{\alpha\beta} \\ & + b_3 K_{\alpha\gamma} K_{\alpha\beta} K_{\beta\gamma} + b_4 K_{\alpha\alpha} K_{\beta\gamma} K_{\beta\gamma} \\ & + b_5 K^\alpha_\alpha K^\beta_\beta K^\gamma_\gamma - b_6 H_{\alpha\delta\epsilon} H_{\beta\delta\epsilon} K^\gamma_\gamma n^\alpha n^\beta + \\ & + b_7 H_{\alpha\gamma}^\delta H_{\beta\delta\epsilon} \gamma\delta\alpha\beta \quad \alpha\beta \quad K \quad n n + b_8 K \quad R_{\alpha\beta} \\ & + b_9 K_{\gamma\nu} n\alpha n\beta R_{\alpha\beta} + b_{10} K_{\alpha\alpha} R \\ & + b_{11} K_{\gamma\delta} n\alpha n\beta R_{\alpha\beta\delta} \\ & + b_{12} H_{\beta\gamma\delta} n\alpha \nabla_\alpha H_{\beta\gamma\delta} \\ & + b_{13} K_{\beta\gamma} n\alpha \nabla_\alpha K_{\beta\gamma} + b_{14} K_{\beta\beta} n\alpha \nabla_\alpha K_{\gamma\gamma} \\ & + b_{15} n^\alpha \nabla_\alpha R \\ & + b_{16} H_{\beta\gamma\delta} H^{\beta\gamma\delta} n^\alpha \nabla_\alpha \Phi + b_{17} K_{\beta\gamma} \\ & \quad \beta\gamma\alpha \quad K \quad n \quad \nabla_\alpha \Phi + b_{18} K^\beta_\beta K^\gamma_\gamma n^\alpha \nabla_\alpha \Phi + b_{17} K_{\beta\gamma} \\ & + b_{19} H_{\beta\delta\epsilon} H_{\gamma\delta\epsilon} n\alpha n\beta n\gamma \nabla_\alpha \Phi \\ & + b_{20} n\alpha n\beta n\gamma R_{\beta\gamma} \nabla_\alpha \Phi + b_{21} n^\alpha R \nabla_\alpha \Phi + \\ & b_{22} K^\beta_\beta \nabla_\alpha \Phi \nabla^\alpha \Phi \\ & + b_{23} n\alpha n\beta \nabla_\alpha \Phi \nabla_\beta K^\gamma_\gamma \\ & + b_{24} K^\gamma_\gamma n^\alpha n^\beta \nabla_\alpha \Phi \nabla_\beta \Phi + b_{25} n^\alpha n^\beta \nabla_\beta \nabla_\alpha K^\gamma_\gamma \\ & + b_{26} K^{\alpha\beta} \nabla_\beta \nabla_\alpha \Phi \\ & + b_{27} K^\gamma_\gamma n^\alpha n^\beta \nabla_\beta \nabla_\alpha \Phi + b_{28} H_{\alpha\gamma\delta} H_{\beta\gamma\delta} n^\alpha \nabla^\beta \Phi \\ & + b_{29} n^\alpha R_{\alpha\beta} \nabla^\beta \Phi + b_{30} K_{\alpha\beta} \nabla^\alpha \Phi \nabla^\beta \Phi \\ & + b_{31} n^\alpha \nabla_\alpha \Phi \nabla_\beta \Phi \nabla^\beta \Phi + b_{32} n^\alpha \nabla_\beta \nabla_\alpha \Phi \nabla^\beta \Phi \\ & + \delta\epsilon \alpha \beta \gamma \quad + \quad b_{33} H_{\alpha\gamma} \quad n n n \nabla_\gamma H_{\beta\delta\epsilon} \\ & + b_{34} n\alpha n\beta n^\gamma \nabla_\alpha \Phi \nabla_\beta \Phi \nabla_\gamma \Phi \end{aligned}$$

$$\begin{aligned}
& +b_{35}n_{\alpha\beta}n_{\gamma}\nabla_{\alpha}\Phi\nabla_{\gamma}\nabla_{\beta}\Phi \\
& +b_{36}n_{\alpha\beta}n_{\gamma}\nabla_{\gamma}\nabla_{\beta}\nabla_{\alpha}\Phi \\
& +b_{37}n_{\alpha\beta}\nabla_{\beta}K_{\alpha\gamma}\nabla^{\gamma}\Phi \\
& +b_{38}n_{\alpha\beta}n_{\gamma}\nabla_{\delta}\nabla_{\gamma}K_{\alpha\beta}
\end{aligned} \tag{19}$$

$$\begin{aligned}
& = \begin{pmatrix} \bar{g}_{ab} + e^{\varphi} & \\ e^{\varphi} g_b & e^{\varphi} \end{pmatrix} \begin{matrix} g_a g_b e^{\varphi} g_a \\ G_{\mu\nu}, \end{matrix} \\
B_{\mu\nu} & = \begin{pmatrix} \bar{b}_{ab} + \frac{1}{2} b_a g_b - \frac{1}{2} & \\ -b_b & b b g_a b^0 \end{pmatrix}, \quad \Phi = \bar{\phi} + \frac{\varphi}{4}
\end{aligned} \tag{20}$$

where g^{-ab} is the metric, b^{-ab} is the B-field, $\bar{\phi}$ is dilaton and g_a, b_b are two vectors in the base space. The reduction of the unit vector n^{μ} is

$$n^{\mu} = \begin{pmatrix} n^a \\ 0 \end{pmatrix} \tag{21}$$

where n^a is the unit vector to the boundary in the base space. Using these reductions, then one reduces the effective actions $S_{\text{eff}} + \partial S_{\text{eff}}$ on the circle to find $S_{\text{eff}}(\psi) + \partial S_{\text{eff}}(\psi)$ where ψ represents all the massless fields in the base space. The Z_2 symmetry then constrains the effective action to satisfy the relation (4). The Z_2 -transformation of the base space fields in (4) are the Buscher rules [23,24] and their higher derivative corrections, i.e.,

$$\psi' = \sum_{m=0}^{\infty} \alpha'^m \psi_m \tag{22}$$

where ψ'_0 represents the Buscher rules. In terms of the reductions (20), they are

$$\varphi = -\varphi, \quad g_a = b_a, \quad b_a = g_a \tag{23}$$

The base space metric, dilaton, b^{-} -field and the unit vector n^a are invariant. The ψ'_0 in (22) represents two derivative corrections to the Buscher rules, and so on.

The corrections to the Buscher rules depend on the scheme that one uses for the gauge invariant couplings [11], and vis versa. Since we have not used the field redefinitions to write the gauge invariant couplings, we are free to consider a specific geometry for imposing the $O(1,1)$ symmetry. For the geometry that we have considered, the unit normal vector n^a on the boundary is independent of the circle on which the T-duality is imposed, i.e., $g^{-ab} n^a n^b = 1$ is invariant under the T-duality. Hence, the base space metric must be invariant under the T-duality at any order of α . On the other hand,

under the reduction (20), the density $e^{-2\Phi} \sqrt{-G}$

reduces to $e^{-2\Phi} \sqrt{-g}$ which may be invariant under the T-duality at any order of α . Therefore, the base space dilaton may be invariant at any order of α . So, for the background that we consider, the base space unit normal vector n^a , the

where b_1, \dots, b_{38} are background independent boundary parameters which may be fixed by imposing the Z_2 symmetry, by the least action principle and by the cosmological constraint (7). Note that the above boundary couplings do have terms with two and three derivatives, hence,

in extremizing the above Lagrangian one faces with the variation of the second and third derivatives of the massless fields on the boundary which are non-zero. As a result, the least action principle constrains the parameters b_1, \dots, b_{38} . In other words, the above boundary action satisfies $\delta(\partial S_1) = 0$ for some specific values of the parameters which, as we will see, are consistent with the T-duality. Note that, as we will see, if one uses the boundary condition that only the massless fields are arbitrary on the boundary, i.e., only the variation of massless fields on the boundary are zero, then the least action principle would constrain more strongly the parameters b_1, \dots, b_{38} which would not be consistent with the Z_2 -symmetry. In the next section we first find the relations between the bulk parameters and the relations between the boundary and bulk parameters by imposing the Z_2 symmetry, and in the section after we impose the least action principle to further constrain the parameters in the boundary action.

3 Z_2 -invariance constraint

Using the assumption that the effective actions in the string theory are independent of the geometry of the spacetime, we now explicitly impose the Z_2 -symmetry on the effective actions to find some relations between the parameters in the gauge invariant couplings (12) and (19). To this end, we consider a particular geometry that $\partial M^{(D)} = \partial M^{(D-1)} \times S^{(1)}$ its bulk and boundary have the structures $M^{(D)}$ and $M^{(D-1)} \times S^{(1)}$.

The manifold $M^{(D)}$ has coordinates $x = (x, y)$ and its boundary $\partial M^{(D)}$ has coordinates $\sigma^{\mu} = (\sigma^a, y)$ where y is the coordinate of the circle $S^{(1)}$. The Kaluza-Klein reduction of the metric and the reduction of B-field and dilaton are [7]

dilaton ϕ^- and the metric g^-_{ab} do not appear in the T-duality transformations. The base space b^- -field which is invariant under the Buscher rules, however, appears in the higher derivative corrections because this field and the vectors g_a, b_a should satisfy a Bianchi identity [29]. Hence, we consider the Tduality transformations that involve only the derivatives of the base space fields ϕ, g_a, b_a, b^-_{ab} .

In order the T-duality constraint (4) to be satisfied, one should add some total derivative terms at the boundary which

are zero by the Stokes's theorem, i.e., $T \equiv -\frac{2}{\kappa^2} \int_{\partial M^{(D-1)}} d^{D-2} \sigma \sqrt{|g|} n_a \nabla_b (e^{-2\tilde{\phi}} F^{ab}) = 0$ (24)

where $F^{ab}(\psi)$ is an arbitrary antisymmetric tensor constructed from the gauge invariant base space fields. It can be α -expanded as $F^{abm} = \sum_{m=0}^{\infty} \alpha^m F_m^{ab}$, which produces an α -expansion for $T = \sum_{m=0}^{\infty} \alpha^m T_m$. Then one can study the Z_2 -constraint (4) at each order of α . Replacing the expansions (8) and (22) in the Z_2 -symmetry (4), one finds the following relation at order α^0 :

$$S_0(\psi) + \partial S_0(\psi) = S_0(\psi'_0) + \partial S_0(\psi'_0) \tag{25}$$

Note that it is impossible to construct the antisymmetric tensor F_0^{ab} at zero derivative order. Hence there is not total derivative on the boundary in this case. This constraint has been used in [25] to find the effective action (5) at order α^0 .

Using the relation (25), one finds that the constraint (4) produces the following relation at order α :

$$S_1(\psi) + \partial S_1(\psi) + T_1(\psi) = S_1(\psi'_0) + \partial S_1(\psi'_0) + \Delta S_0 + \Delta \partial S_0 \tag{26}$$

where $T_1(\psi)$ is the total derivative terms (24) at order α , and $\Delta S_0, \Delta \partial S_0$ are defined in the following α -expansions:

$$\begin{aligned} S_0(\psi'_0 + \alpha' \psi'_1) - S_0(\psi'_0) &= \alpha' \Delta S_0 + \dots \\ \partial S_0(\psi'_0 + \alpha' \psi'_1) - \partial S_0(\psi'_0) &= \alpha' \Delta \partial S_0 + \dots \end{aligned} \tag{27}$$

where dots represent some terms at higher orders of α in which we are not interested in this paper. The constraint (26) can be written as

$$\begin{aligned} S_1(\psi) - S_1(\psi'_0) - \Delta S_0 &= -T_1(\psi) \\ -(\partial S_1(\psi) - \partial S_1(\psi'_0) - \Delta \partial S_0) & \end{aligned} \tag{28}$$

The terms on the left-hand side are in the $(D-1)$ -dimensional base space whereas the terms on the right-hand side are in its boundary. Only the total derivative terms in the $(D-1)$ -

dimensional action can contribute to its boundary action. Hence, the bulk actions on the left-hand side should be some total derivative terms, i.e.,

$$S_1(\psi) - S_1(\psi'_0) - \Delta S_0 = \frac{2}{\kappa^2} \int_{M^{(D-1)}} d^{D-1} x \sqrt{-\tilde{g}} \nabla_a (A_1^a e^{-2\tilde{\phi}}) \tag{29}$$

where A_1^a is a vector made of the covariant derivative of the massless fields in the base space at order α . The total derivative terms then produce some boundary terms using the Stokes's theorem.

The Stokes's theorem in the base space is

$$\begin{aligned} \int_{M^{(D-1)}} d^{D-1} x \sqrt{-\tilde{g}} \nabla_a (A_1^a e^{-2\tilde{\phi}}) \\ = \int_{\partial M^{(D-1)}} d^D \dots -2\sigma \sqrt{|\tilde{g}|} n_a A_1^a | \dots e^{-2\tilde{\phi}} \end{aligned} \tag{30}$$

where n^a is the unit vector orthogonal to the boundary in the base space and the boundary is specified by the functions $x^a = x^a(\sigma^a)$. The unit vector is outward-going (inwardgoing) if the boundary is spacelike (timelike). The metric in the square root on the right-hand side is the induced metric, i.e.,

$$\begin{aligned} \partial x_a \partial x_b \\ g^a b^- = \partial \sigma \dots \dots a^- \partial \sigma b^- g^-_{ab} \end{aligned} \tag{31}$$

Using the Stokes's theorem to transfer the bulk total derivative terms on the right-hand side of (29) to the boundary, then the Z_2 -symmetry on the effective action at order α produces the bulk constraint (29) as well as the following constraint on the boundary couplings:

$$\begin{aligned} \partial S_1(\psi) - \partial S_1(\psi'_0) - \Delta \partial S_0 + T_1(\psi) + \frac{2}{\kappa^2} \\ \times \int_{\partial M} d^D \dots -2\sigma \sqrt{|\tilde{g}|} n_a A_1^a e^{-2\tilde{\phi}} = \dots \end{aligned} \tag{32}$$

where A_1^a has to be found from the bulk constraint (29). The constraint (29) and (32) produce some relations between the parameters in the gauge invariant couplings (12) and (19).

3.1 Bulk constraint

The bulk constraint (29) has been used in [11] to find some relations between the parameters in the action (12) for the

most general T-duality transformation at order α . The total derivative terms have been ignored in calculating ΔS_0 in [11] because in that calculation it was assumed spacetime has no boundary. Hence, the result in [11] can not be used for the present case that there is boundary. In the present case all total derivative terms in the base space should be taken into account. So we solve the constraint (29) in this section to keep track of the total derivative terms carefully.

To find ΔS_0 from the expansion (27), one needs the reduction of the bulk action at order α^0 . Using the reductions (20), one finds the reduction of the bulk action S_0 to be

$$S_0(\psi) = -\frac{2}{\kappa^2} \int d^{D-1}x e^{-2\bar{\phi}} \sqrt{-\bar{g}} \times \left[\bar{R} - \nabla^a \nabla_a \varphi - \frac{1}{4} \nabla_a \varphi \nabla^a \varphi - \frac{1}{4} (e^\varphi V^2 + e^{-\varphi} W^2) + 4 \nabla_a \bar{\phi} \nabla^a \bar{\phi} + 2 \nabla_a \bar{\phi} \nabla^a \varphi - \frac{1}{12} \bar{H}_{abc} \bar{H}^{abc} \right] \quad (33)$$

where V_{ab} is field strength of the $U(1)$ gauge field g_a , i.e., $V_{ab} = \partial_a g_b - \partial_b g_a$, and W_{uv} is field strength of the $U(1)$ gauge field b_a , i.e., $W_{ab} = \partial_a b_v - \partial_b b_a$. The three-form H^- is defined as $H^-_{abc} = \hat{H}^-_{abc} - \frac{3}{2} g_{[a} W_{bc]} - \frac{3}{2} b_{[a} V_{bc]}$ where the three-form \hat{H}^- is field strength of the two-form b^-_{ab} in (20). Since H^- is not exterior derivative of a two-form, it satisfies anomalous Bianchi identity, whereas the W, V satisfy the ordinary Bianchi identity, i.e.,

$$\begin{aligned} \partial_{[a} \bar{H}_{bcd]} &= -\frac{3}{2} V_{[ab} W_{cd]} \\ \partial_{[a} W_{bc]} &= 0 \\ \partial_{[a} V_{bc]} &= 0 \end{aligned} \quad (34)$$

Our notation for making antisymmetry is such that e.g., $g_{[a} W_{bc]} = \frac{1}{3} (g_a W_{bc} - g_b W_{ac} - g_c W_{ba})$.

The Z_2 -transformations at order α^0 , i.e., ψ'_i , are given by the Buscher rules (23), and at order α are given by the following: $\varphi = -\varphi + \alpha' \Delta \varphi$, $g'_a = b_a + \alpha' e^{\varphi/2} \Delta g_a$, $b'_a = g_a + \alpha' e^{-\varphi/2} \Delta b_a$, $\bar{H}'_{abc} = \bar{H}_{abc} + \alpha' \Delta \bar{H}_{abc}$ (35) where the corrections $\Delta \varphi, \Delta g_a, \Delta b_a$ contain some contractions of $\nabla \varphi, e^{\varphi/2} V, e^{-\varphi/2} W, H^-$ at order α . Note that the base space metric, dilaton, b^- -field and the unit vector n^a remain invariant at order α . Since the transformations must

form the Z_2 -group, the corrections satisfy the following relations [11]:

$$\begin{aligned} -\Delta \varphi(\psi) + \Delta \varphi(\psi'_0) &= 0 \\ \Delta b_a(\psi) + \Delta g_a(\psi'_0) &= 0 \\ \Delta g_a(\psi) + \Delta b_a(\psi'_0) &= 0 \\ \Delta \bar{H}_{abc}(\psi) + \Delta \bar{H}_{abc}(\psi'_0) &= 0 \end{aligned} \quad (36)$$

Then the corrections should have the following terms:

$$\begin{aligned} \Delta \varphi &= e_3 \partial_a \varphi \partial^a \varphi + e_1 \bar{H}^2 + e_2 (e^\varphi V^2 + e^{-\varphi} W^2) \\ g_a &= e_4 e^{\varphi/2} H^-_{abc} V_{bc} + e_5 e^{-\varphi/2} \partial_b \varphi W_{ab} \\ \Delta b_a &= -e_4 e^{-\varphi/2} \bar{H}_{abc} W^{bc} + e_5 e^{\varphi/2} \partial^b \varphi V_{ab} \end{aligned} \quad (37)$$

where e_1, \dots, e_5 are some parameters that should be fixed by the Z_2 -symmetry of the effective actions. Note that we did not include the corrections which depend on the derivative of the base space dilaton and metric. The correction ΔH_{abc} is related to the corrections $\Delta g_a, \Delta b_a$ through the following relation which is resulted from the Bianchi identity (34):

$$\Delta \bar{H}_{abc} = H^-_{abc} - 3e^{-\varphi/2} W_{[ab} \Delta b_{c]} - 3e^{\varphi/2} \Delta g_{[a} V_{bc]} \quad (38)$$

where H^-_{abc} is a $U(1) \times U(1)$ gauge invariant closed 3-form at order α which is odd under parity. It has the following terms:

$$H^-_{abc} = e_6 \partial_{[a} W_{b^d} V_{c]d} + e_7 \partial_{[a} H^-_{bc]d} \nabla^d \varphi \quad (39)$$

where e_6, e_7 are two other parameters that the Z_2 -symmetry of the effective action should fix them.

Using the reduction (33), then one can calculate ΔS_0 from the expansion (27) in terms of above corrections, i.e.,

$$\begin{aligned} S_0 &= -\frac{2}{\kappa^2} \int d^{D-1}x e^{-2\bar{\phi}} \sqrt{-\bar{g}} \left[\frac{1}{4} e^\varphi V^2 - e^{-\varphi} W^2 - \varphi \right. \\ &\quad + \frac{1}{2} e^{-\varphi/2} \partial_b \varphi W_{ab} g_a - 1 e^{\varphi/2} \partial_b \varphi V_{ab} b_a \\ &\quad - \frac{1}{6} \bar{H}^{abc} \Delta \bar{H}_{abc} \\ &\quad + \frac{1}{2} (\partial_a \varphi + 4 \partial_a \bar{\phi}) \nabla^a (\Delta \varphi) - \nabla_a \nabla^a (\Delta \varphi) \\ &\quad \left. + e^{-\varphi/2} W_{ab} \nabla^b (\Delta g^a) + e^{\varphi/2} V_{ab} \nabla^b (\Delta b^a) \right] \quad (40) \end{aligned}$$

where no integration by part has been used. One can check that up to some total derivative terms the above expression become the same as the corresponding expression in [11] for $\Delta \bar{\phi} = \Delta \bar{g}_{ab} = 0$ (i.e., eq.(23) in [11]), in which the total derivative terms were ignored. Note that ΔS_0 is not even or odd under the Buscher rules. However, up to some total derivative terms it is odd under the Buscher rules [11].

Using the reductions (20), it is straightforward to reduce the effective action S_1 with the Lagrangian density (12) to find $S_1(\psi)$, and then calculate its transformation under the Buscher rules (23), i.e., $S_1(\psi'_0)$. See [11], for the details of this calculation. Note that $S_1(\psi) - S_1(\psi'_0)$ is odd under the Buscher rules, however, since ΔS_0 is not even or odd under the Buscher rules, the vector $A_1^a(\psi)$ in (29) is not even or odd under the Buscher rules either. Hence, the vector $A_1^a(\psi)$ should contain all even-parity contractions of $\partial\phi$, $\partial\phi^-$, $e^{\varphi/2}V, e^{-\varphi/2}W, H^-, R^-$ and their derivatives at order α with unknown coefficients. Using the package "xAct", one finds it has 48 terms, i.e.,

$$\begin{aligned}
 A_1^a = & j_1 \bar{H}^{bcd} V_{bc} W_{ad} + j_2 H^{-acd} V_{bc} W_{bd} + j_3 H^{-bcd} V_{ab} W_{cd} \\
 & + j_4 H^{-bcd} \nabla_a H^{-bcd} + j_7 \nabla_a R \\
 & + \\
 & + e^\varphi j_{17} V^{bc} \nabla^a V_{bc} + e^{-\varphi} j_{20} W^{bc} \nabla \\
 & + j_9 \bar{H}^{bcd} \bar{H}^{bcd} \nabla^a \bar{\phi} + j_{11} R \nabla^a \bar{\phi} + e^\varphi j_{13} \\
 & e^{-\varphi} j_{15} W_{bc} W^{bc} \nabla^a \phi \\
 & V_{bc} V_{bc} \nabla_a \phi^- \\
 & W_{bc} \\
 & + j_{23} H^{-bcd} H^{-bcd} \nabla_a \phi + j_{25} R \nabla_a \phi \\
 & + j_8 \nabla_b R^{ab} + j_{41} \nabla^a \bar{\phi} \nabla_b \nabla^b \bar{\phi} + j_{43} \nabla^a \phi \nabla_b \nabla^b \bar{\phi} \\
 & + j_{31} \nabla^a \nabla_b \nabla^b \bar{\phi} + j_{45} \nabla^a \bar{\phi} \nabla_b \nabla^b \phi + j_{47} \nabla^a \phi \nabla_b \nabla^b \phi \\
 & + j_{33} \nabla^a \nabla_b \nabla^b \phi + e^\varphi j_{27} V_{bc} V_{bc} \nabla_a \phi + e^{-\varphi} \\
 & j_{29} W_{bc} W_{bc} \nabla_a \phi \\
 & + j_{32} \nabla_b \nabla^b \nabla^a \bar{\phi} + j_{34} \nabla_b \nabla^b \nabla^a \phi + j_{10} \bar{H}^{acd} \bar{H}^{bcd} \nabla^b \bar{\phi} \\
 & + j_{12} R \\
 & + e^{-\varphi} j_{16} W^{ac} W_{bc} \nabla^b \bar{\phi} + j_{35} \nabla^a \bar{\phi} \nabla_b \bar{\phi} \nabla^b \bar{\phi} \\
 & + j_{36} \nabla^a \phi \nabla_b \bar{\phi} \nabla^b \bar{\phi} + j_{37} \nabla^a \bar{\phi} \nabla_b \phi \nabla^b \bar{\phi} \\
 & + j_{38} \nabla^a \phi \nabla_b \phi \nabla^b \bar{\phi} + j_{42} \nabla_b \nabla^a \bar{\phi} \nabla^b \bar{\phi} \\
 & + j_{46} \nabla_b \nabla^a \phi \nabla^b \bar{\phi} \\
 & + j_{24} \bar{H}^{acd} \bar{H}^{bcd} \nabla^b \phi + j_{26} R_{ab} \nabla^b \phi \\
 & + e^\varphi j_{28} V_{ac} V_{bc} \nabla^b \phi + e^{-\varphi} j_{30} W_{ac} W_{bc} \nabla^b \phi \\
 & + j_{39} \nabla_a \phi^- \nabla_b \phi \nabla^b \phi + j_{40} \nabla_a \phi \nabla^b \phi \nabla^b \phi \\
 & + j_{44} \nabla_b \nabla_a \phi^- \nabla^b \phi + j_{48} \nabla_b \nabla_a \phi \nabla^b \phi \\
 & + e^\varphi j_{18} V_{bc} \nabla_c V_{ab} + e^\varphi j_{19} V_{ab} \nabla_c V_{bc} \\
 & + e^{-\varphi} j_{21} W_{bc} \nabla_c W_{ab} + e^{-\varphi} j_{22} W_{ab} \nabla_c W_{bc}
 \end{aligned}$$

$$+ j_5 H^{-bcd} \nabla_d H^{-abc} + j_6 H^{-abc} \nabla_d H^{-bcd} \tag{41}$$

where j_1, \dots, j_{48} are 48 parameters. Note that some of the above terms are related by the Bianchi identities. We will remove them after imposing the constraint (29).

To find the parameters which satisfy the bulk constraint (29), one should write the covariant derivatives and the curvatures in (29) in terms of partial derivative of the base space

metric, and write the partial derivatives of the base space field strength H^-, W, V in terms of the potentials b^{-ab}, g_a, b_a and then goes to the local frame in which the first derivative of the metric is zero [22]. The coefficients of the resulting independent terms in the local frame then should be zero. This gives some linear equations involving all the parameters. We find the following solution in terms of the parameter a_1, a_{12} :

$$\begin{aligned}
 a_{10} = & -16a_1, a_{11} = 0, a_{13} = 384a_1 - 12a_{12}, a_{14} = 768a_1 - \\
 & 24a_{12}, a_{15} = 96a_1 - 6a_{12}, a_{16} = -768a_1 + 24a_{12}, a_{17} = -768a_1 \\
 & + 24a_{12}, a_{18} = 48a_1, a_{19} = 0, a_2 = -9a_1 + (3a_{12})/8, a_{20} = -24a_1 \\
 & + (3a_{12})/2, a_3 = a_1/3 - a_{12}/96, a_4 = 72a_1 - 3a_{12}, a_5 = -192a_1 \\
 & + 6a_{12}, a_6 = -8a_1 + a_{12}/4, a_7 = 48a_1 - (3a_{12})/2, a_8 = 24a_1, \\
 & a_9 = -12a_1, e_6 = -288a_1, e_7 = 0, e_1 = 0, e_2 = 24a_1, e_3 = \\
 & 48a_1, e_4 = 24a_1, e_5 = 48a_1, j_1 = -24a_1, j_{10} = 0, j_{11} = 0, \\
 & j_{12} = 0, j_{13} = -48a_1, j_{14} = 96a_1, j_{15} = 48a_1, j_{16} = -96a_1, j_{17} \\
 & = -48a_1, j_{18} = 48a_1, j_{19} = -48a_1, j_2 = 0, j_{20} = -48a_1, j_{21} = \\
 & -48a_1, j_{22} = 48a_1, j_{23} = -8a_1 + a_{12}/2, j_{24} = 48a_1 - 3a_{12}, \\
 & j_{25} = 192a_1 - 6a_{12}, j_{26} = -384a_1 + 12a_{12}, j_{27} = -72a_1 + \\
 & (3a_{12})/2, \\
 & j_{28} = 192a_1 - 6a_{12}, j_{29} = -24a_1 + (3a_{12})/2, j_3 = 24a_1, j_{30} = 192a_1 \\
 & - 6a_{12}, j_{31} = 0, j_{32} = 0, j_{33} = 0, j_{34} = 0, j_{35} = 0, j_{36} = 384a_1 - \\
 & 12a_{12}, j_{37} = -768a_1 + 24a_{12}, j_{38} = 0, j_{39} = 0, j_4 = 0, j_{40} = 24a_1 - \\
 & (3a_{12})/4, j_{41} = 0, j_{42} = 0, j_{43} = 0, j_{44} = 0, j_{45} = 0, j_{46} = 0, j_{47} = 0, \\
 & j_{48} = -96a_1, j_5 = 0, j_6 = 0, j_8 = -2j_7, j_9 = 0
 \end{aligned}$$

(42)

When replacing them into (37) and (39), one finds the following corrections to the Buscher rules:

$$\begin{aligned}\Delta\varphi &= 24a_1\left(2\partial_a\varphi\partial^a\varphi + e^\varphi V^2 + e^{-\varphi}W^2\right) \\ \Delta g_a &= 24a_1\left(2e^{-\varphi/2}\partial^b\varphi W_{ab} + e^{\varphi/2}\bar{H}_{abc}V^{bc}\right) \\ \Delta b_a &= 24a_1\left(2e^{\varphi/2}\partial^b\varphi V_{ab} - e^{-\varphi/2}\bar{H}_{abc}W^{bc}\right)\end{aligned}$$

$$H^-{}_{abc} = -288a_1\partial_{[a}(W_b{}^dV_{c]d}) - 3e^{\varphi/2}V_{1ab}\Delta g_{c1} - 3e^{-\varphi/2}W_{1ab}\Delta b_{c1}(43)$$

These transformations are those have been found in [29]² for $24a_1 = -\lambda_0$.

Since the above corrections are independent of the parameter a_{12} , the solution (42) produces two multiplets. One with the overall coefficient a_1 which is invariant under the Buscher rules plus the above higher derivative corrections, and the other one with the overall coefficient a_{12} which is invariant under the Buscher rules. The T-duality then can not fix a relation between these two parameters. We find the relation between them by using the fact that the sphere-level S-matrix elements of massless vertex operators in the string theory have simple poles reflecting the standard propagators for the massless and massive fields in the amplitude. At the low energy, one expands the massive propagators to find an amplitude in terms of only massless fields. The massless poles at the low energy are still simple poles. They should be reproduced by effective actions which have standard massless propagators. The last term in the effective action (12) changes the standard propagators of the B -field. However, for the following relation between a_1 and a_{12} :

$$a_{12} = 16a_1 \quad (44)$$

the coefficient of this term become zero after imposing the T-duality constraint (42), i.e., $a_{20} = 0$. Interestingly, imposing this relation, one finds the curvature terms in the action (12) also becomes proportional to the Gauss–Bonnet gravity which does not change the standard propagator of the metric.

Imposing the relation (44) and the Z_2 -constraint (42), one finds the following bulk action:

$$\begin{aligned}\mathcal{L}_1 &= 24a_1\left[\frac{1}{24}H_\alpha{}^{\delta\epsilon}H^{\alpha\beta\gamma}H_{\beta\delta}{}^\epsilon H_{\gamma\epsilon}{}^\delta\right. \\ &\quad -\frac{1}{8}H_{\alpha\beta}{}^\delta H^{\alpha\beta\gamma}H_\gamma{}^{\epsilon\delta}H_{\delta\epsilon}{}^\gamma \\ &\quad +\frac{1}{144}H_{\alpha\beta\gamma}H^{\alpha\beta\gamma}H_{\delta\epsilon\epsilon}H^{\delta\epsilon\epsilon} \\ &\quad +H_\alpha{}^{\gamma\delta}H_{\beta\gamma\delta}R^{\alpha\beta} - 4R_{\alpha\beta}R^{\alpha\beta} \\ &\quad -\frac{1}{6}H_{\alpha\beta\gamma}H^{\alpha\beta\gamma}R + R^2 + R_{\alpha\beta\gamma\delta}R^{\alpha\beta\gamma\delta} \\ &\quad -\frac{1}{2}H_\alpha{}^{\delta\epsilon}H^{\alpha\beta\gamma}R_{\beta\gamma\delta\epsilon} \\ &\quad -\frac{2}{3}H_{\beta\gamma\delta}H^{\beta\gamma\delta}\nabla_\alpha\nabla^\alpha\Phi \\ &\quad +\frac{2}{3}H_{\beta\gamma\delta}H^{\beta\gamma\delta}\nabla_\alpha\Phi\nabla^\alpha\Phi + 8R\nabla_\alpha\Phi\nabla^\alpha\Phi \\ &\quad +16\nabla_\alpha\Phi\nabla^\alpha\Phi\nabla_\beta\nabla^\beta\Phi \\ &\quad -16R_{\alpha\beta}\nabla^\alpha\Phi\nabla^\beta\Phi - 16\nabla_\alpha\Phi\nabla^\alpha\Phi\nabla_\beta\Phi\nabla^\beta\Phi \\ &\quad \left.+2H_\alpha{}^{\gamma\delta}H_{\beta\gamma\delta}\nabla^\beta\nabla^\alpha\Phi\right] \quad (45)\end{aligned}$$

Since all background independent parameters in the gauge invariant action (12) are fixed in the particular geometry which has one circle, then the above action should be valid for any other geometry as well, e.g., if one considers a geometry which has a torus T^2 , then the reduction of the above action should have the symmetry $O(2,2)$. The above action is exactly the off-shell Lagrangian that has been found in [8] by imposing various field redefinitions on the on-shell action in the minimal scheme.

The relation (44) and the Z_2 -constraint (42) produces the total derivative terms in (29) that their corresponding vector A^a_1 is the following:

$$\begin{aligned}A^a_1 &= 24a_1\left[\bar{H}_{bcd}V_{ab}W_{cd} - H^-{}_{bcd}V_{bc}W_{ad}\right. \\ &\quad -2e_\varphi V_{bc}V_{bc}\nabla^a\phi^- \\ &\quad +2e^{-\varphi} \\ &\quad \left.{}^\varphi_{\varphi ac}W^{bc}\nabla^a W_{bc} + 4R\nabla^a\varphi - \right. \\ &\quad \left. W_{bc}W_{bc}\nabla^a\phi^- - 2e_\varphi V_{bc}\nabla^a V_{bc}\right. \\ &\quad \left. +2e_\varphi V_{bc}V_{bc}\nabla^a\varphi\right. \\ &\quad \left. +8\nabla^a\varphi\nabla_b\bar{\phi}\nabla^b\bar{\phi} - 16\nabla^a\bar{\phi}\nabla_b\varphi\nabla^b\bar{\phi} - \right. \\ &\quad \left. \varphi_{\varphi ac} \quad b \quad \varphi_{\varphi ac} \quad b\right. \\ &\quad \left. +4e V_{bc}\nabla^a\varphi\right.\end{aligned}$$

² The sign of the first term in the last line of (43), however, is different than the one appears in [29] which is a typo [30].

$$\begin{aligned}
 & \phi^- - 4e^- W_{bc} \nabla \phi^- \\
 & \quad + 4e^- V_{bc} \nabla \phi + 4e^- W_{bc} \nabla \phi \\
 & \quad + \frac{1}{2} \nabla^a \phi \nabla_b \phi \nabla^b \phi - 4 \nabla_b \nabla^a \phi \nabla^b \phi + 2e^\varphi V^{bc} \nabla_c V_{ab} \\
 & \quad - 2e\varphi V_{ab} \nabla_c V_{bc} - 2e^{-\varphi} W_{bc} \nabla_c W_{ab} \\
 & \quad + 2e^{-\varphi} W^{ab} \nabla_c W_b^c \Big] \tag{46}
 \end{aligned}$$

Note that the above vector is not odd or even under the Buscher rules.

The presence of the total derivative terms indicates that the bulk action alone is not invariant under the Z_2 transformations. Using the Stokes’s theorem, these anomalous terms are transferred to the boundary and then should be cancelled with the anomalous terms in the boundary action. The anomalous terms from the bulk action are then the following terms:

$$\begin{aligned}
 & \frac{dD}{dM} \left[2\sigma \sqrt{|g|} n_a A_1^a e^{2\phi} \Big| \right] \\
 & = 24a_1 \int_{\partial M} d^{D-2} \sigma \sqrt{|g|} e^{-2\phi} n_a \left[\bar{H}_{bcd} V_{ab} W_{cd} \right. \\
 & \quad - H^{-bcd} V_{bc} W_{ad} - 2e^{-\varphi} V_{bc} V_{bc} \nabla_a \phi^- \\
 & \quad + 2e^{-\varphi} W_{bc} W_{bc} \nabla_a \phi^- - 2e\varphi V_{bc} \nabla_a V_{bc} \\
 & \quad - 2e^{-\varphi} W_{bc} \nabla_a W_{bc} + 4R \nabla_a \phi \\
 & \quad - 2e\varphi V_{bc} V_{bc} \nabla_a \phi + 4e\varphi V_{ac} V_{bc} \nabla_b \phi^- \\
 & \quad - 4e^{-\varphi} W^{ac} W_{bc} \nabla^b \bar{\phi} \\
 & \quad + 8 \nabla^a \phi \nabla_b \bar{\phi} \nabla^b \bar{\phi} - 16 \nabla^a \phi \nabla_b \phi \nabla^b \phi \\
 & \quad - 8R_b \nabla \phi + 4e^- V_{bc} \nabla_b \phi + 4e^{-\varphi} W_{ac} W_{bc} \nabla_b \phi \\
 & \quad + \frac{1}{2} \nabla^a \phi \nabla_b \phi \nabla^b \phi - 4 \nabla_b \nabla^a \phi \nabla^b \phi \\
 & \quad + 2e\varphi V_{bc} \nabla_c V_{ab} - 2e\varphi V_{ab} \nabla_c V_{bc} \\
 & \quad \left. - 2e^{-\varphi} W^{bc} \nabla_c W_b^a + 2e^{-\varphi} W^{ab} \nabla_c W_b^c \right] \tag{47}
 \end{aligned}$$

In the next subsection we study the Z_2 -transformations of the boundary action (19).

3.2 Boundary constraint

To impose the Z_2 -constraint (32) on the boundary terms, one first assumes that the boundary is specified by the functions $x^\mu(\sigma^{\tilde{a}}) = (x^a(\sigma^{\tilde{a}}, y))$ where the functions $x^a(\sigma^{\tilde{a}})$ represent the boundary in the base space which are independent of the y -coordinate. Then the reduction of the induce metric (9) becomes

$$\begin{aligned}
 & = \left(\partial \sigma^{\tilde{a}} \partial x^{\tilde{a}} \partial x^{\tilde{b}} g^{-ab} + e^{\int \varphi} \frac{g_a g_b}{\partial \sigma^{\tilde{a}}} \frac{\partial x^{\tilde{a}}}{\partial \sigma^{\tilde{a}}} (e^\varphi g_a) \right) \\
 & g_{\mu\nu} \left(\frac{x}{\partial \sigma^{\tilde{a}}} \frac{\varphi}{\partial \sigma} \partial(b(e g b)) - e\varphi \right) \tag{48}
 \end{aligned}$$

Using this and the reduction of dilaton in (20), one finds the following reduction:

$$e^{-2\phi} \sqrt{|g|} = e^{-2\phi} \sqrt{|g|} \tag{49}$$

where g is the determinate of the induced metric (31). For the background that we have considered, it is invariant under the Z_2 -transformations to all orders of α .

Using the reduction (21) for the normal vector and the reduction (20) for metric, one finds the reduction of the extrinsic curvature to be

$$G^{\mu\nu} K_{\mu\nu} = \bar{g}^{ab} \bar{K}_{ab} + \frac{1}{2} n^a \nabla_a \varphi \tag{50}$$

where K_{ab} is the extrinsic curvature of the boundary of the base space. So the reduction of the boundary action ∂S_0 in

$$\partial S_0(\psi) = -\frac{4}{\kappa^2} \int d^{D-2} \sigma e^{-2\phi} \sqrt{|\tilde{g}|} \left[\bar{g}^{ab} \bar{K}_{ab} + \frac{1}{2} n^a \nabla_a \varphi \right] \tag{51}$$

which is invariant under $U(1) \times U(1)$ gauge transformations.

In using the boundary constraint (32), one needs to calculate $\Delta \partial S_0$. Using the above reduction, one can calculate $\Delta \partial S_0$ in terms of the T-duality transformations (35), i.e.,

$$\Delta \partial S_0 = -\frac{4}{\kappa^2} \int d^{D-2} \sigma e^{-2\phi} \sqrt{|\tilde{g}|} \left[\frac{1}{2} n^a \nabla_a (\Delta \varphi) \right] \tag{52}$$

where φ is given in (43). Note that in finding the above result we have used the fact that in the background that we have considered, the unit vector n^a and the base space metric and dilaton are invariant under the Z_2 -transformations.

Using the reductions (20), it is straightforward to reduce the effective action ∂S_1 with the Lagrangian density (19) to find $\partial S_1(\psi)$, and then calculate its transformation under the Buscher rules (23), i.e., $\partial S_1(\psi'_0)$. See [13,26], for the reduction of different tensors in the boundary Lagrangian (19). Note that for simplicity of calculations in the boundary, we assume the metric of the base space is flat, i.e., $g^{-ab} = \eta_{ab}$. This calculation produces the expression $\partial S_1(\psi) - \partial S_1(\psi'_0)$ in (32). The anomalous couplings from the T-duality of the bulk action is also given in (47). Finally, one needs to add the total derivative terms $T_1(\psi)$ to the constraint (32). According to the Stokes's theorem, the total derivative terms in the boundary which have the following structure are zero:

$$= -\frac{2}{\kappa^2} \int_{\partial} T_1(\psi) \quad d^D \quad (53)$$

where F^{ab} is an arbitrary even-parity antisymmetric tensor constructed from $U(1) \times U(1)$ gauge invariant tensors $n, \partial n, \partial \phi, \partial \varphi, e^{-\varphi/2} W, e^{\varphi/2} V, H^-$ at two derivative order. Using the package "xAct", one can construct this tensor with arbitrary coefficients.

Having calculated all terms in (32), one should then impose the Bianchi identities (34) in the flat base space which can be done by writing the derivatives of the field strengths V, W, H^- in terms of potentials g_a, b_a, b^{-ab} . To impose the identities corresponding to the unit vector n^a in the base space, we also write it in terms of the function f using (13), i.e.,

$$n^a = (\partial_b f \partial^b f)^{-1/2} \partial^a f \quad (54)$$

where we have used the fact that the function f should be independent of the killing coordinate y , i.e., $\partial_y f = 0$. The coefficients of the resulting independent terms then should be zero. They produce some linear equations involving a_1 and the parameters in (19) and in (53).

We find the following non-zero couplings in the boundary:

$$\partial \mathcal{L}_1 = b_1 H_{\beta\gamma\delta} H^{\beta\gamma\delta} K^\alpha{}_\alpha + 24a_1 H_\alpha{}^{\gamma\delta} H_{\beta\gamma\delta} K^{\alpha\beta} - 2$$

$$+ b_{11} K_\alpha{}^\gamma K^{\alpha\beta} K_{\beta\gamma} + (24b_1$$

$$- b_{11}) H_\alpha{}^{\delta\epsilon} H_{\beta\delta\epsilon} K^\gamma n^\alpha n^\beta \quad (55)$$

$$+ 12(-16a_1 + b_{12}) K^{\alpha\beta} R_{\alpha\beta}$$

$$+ (96a_1 + 24b_1 - 12b_{12} - \frac{1}{2}b_{17} + 2b_{19})$$

$$\times K_{\gamma\nu\alpha n\beta} R_{\alpha\beta} - 12b_1 K_{\alpha\alpha} R$$

$$+ b_{11} K_{\gamma\delta n\alpha n\beta} R_{\alpha\beta}$$

$$+ b_{12} H_{\beta\gamma\delta n\alpha} \nabla_\alpha H_{\beta\gamma\delta} - 2b_1 H_{\beta\gamma\delta} H^{\beta\gamma\delta} n^\alpha \nabla_\alpha \Phi$$

$$+ b_{17} K_{\beta\gamma} K_{\beta\nu\alpha} \nabla_\alpha \Phi + b_{18} K_{\beta\gamma} K^\gamma n^\alpha \nabla_\alpha \Phi \quad \beta$$

$$+ b_{19} H_\alpha{}^{\delta\epsilon} H_{\gamma\delta\epsilon} n^\alpha n^\beta \nabla_\gamma \nabla_\nu \Phi \quad \alpha$$

$$+ (-192a_1 + 24b_{12} + b_{17} -$$

$$4b_{19}) n_{\alpha n\beta n^\gamma} R_{\beta\gamma} \nabla_\alpha \Phi$$

$$+ 24(8a_1 + b_1) n^\alpha R \nabla_\alpha \Phi - 48b_1 K_{\beta\gamma} \nabla_\alpha \Phi \nabla^\alpha \Phi_\beta$$

$$- \frac{2}{1} (192a_1 - 48b_1 - \frac{b}{18}) K^\gamma n^\alpha n^\beta \nabla_\alpha \Phi \nabla_\beta \Phi$$

$$+ 24(-16a_1 - 2b_1 +$$

$$b_{12}) K_{\beta\alpha} \nabla^\alpha \nabla^\beta \Phi + 24(-16a_1 - 2b_1 +$$

$$+ 4b_{19}) K^\gamma n^\alpha n^\beta \nabla_\beta \nabla_\alpha \Phi$$

$$+ 48(8a_1 - b_1) n^\alpha R_{\alpha\beta} \nabla^\beta \Phi + \frac{1}{1} \nabla^\alpha \Phi \nabla^\beta \Phi$$

$$- \frac{1}{1} b_{11} - 384a_1 K_{\alpha\beta}$$

$$96(4a_1 - b_1) n^\alpha \nabla_\alpha \Phi \nabla_\beta \Phi \nabla^\beta \Phi \quad (384a_1 + 96b_1 +$$

$$48b_{12} + 2b_{17} - 8b_{19})$$

$$4$$

$$- 6b_{12}) H_\alpha{}^{\delta\epsilon} n^\alpha n^\beta n^\gamma \nabla_\gamma H_{\beta\delta\epsilon}$$

$$\alpha \beta \gamma$$

$$+ \times n n n$$

$$+ b_{38} n_\alpha n_\beta n_\gamma n_\delta \nabla^\alpha \nabla^\beta \nabla_\gamma K_{\alpha\beta}$$

$$4$$

$$+ (-576a_1 - 96b_1 - \frac{b}{18}) n^\alpha n^\beta n^\gamma \nabla_\alpha \nabla_\beta \nabla_\gamma \Phi -$$

$$1 - \quad 1 + 18 \quad \alpha \quad \beta$$

$$\gamma$$

which has the bulk parameter a_1 and 7 boundary parameters $b_1, b_{11}, b_{12}, b_{17}, b_{18}, b_{19}, b_{38}$. The corresponding antisymmetric tensor F^{ab} in the Stokes's theorem is

$$F_{ab} = 6(8a_1 - b_{12}) e_{\phi n b n c} V_{ad} V_{cd}$$

$$- 6(8a_1 - b_{12}) e_{\phi n a n c} V_{bd} V_{cd}$$

$$- 6(8a_1 - b_{12}) e_{\phi n b n c} W_{ad} W_{cd}$$

$$+ (-8b_1 - \frac{1}{6}b_{18}) K^\alpha{}_\alpha K^\beta{}_\beta K^\gamma{}_\gamma - 6(-8a_1 +$$

$$b_{12}) e_{\phi n a n c} W_{bd} W_{cd}$$

$$\begin{aligned}
 &+24b_1n^b\partial^a\varphi\partial_c n^c - 24b_1n^a\partial^b\varphi\partial_c n^c \\
 &-48(8a_1 + b_1)n^b n^c \partial^a \varphi \partial_c \bar{\phi} \\
 &+48(8a_1 + b_1)n^a n^c \partial^b \varphi \partial_c \bar{\phi} \\
 &-24b_1 n b \partial_a n c \partial_c \varphi + 24b_1 n a \partial_b n c \partial_c \varphi
 \end{aligned}$$

which has the bulk parameter a_1 and the boundary parameters b_1, b_{12} . The background independent parameters $b_1, b_{11}, b_{12}, b_{17}, b_{18}, b_{19}, b_{38}$ may be fixed further by considering the geometry which has the tours T^2 in which we are not interested in this paper. In this paper, however, we further constrain the parameters by using the least action principle, and by considering the geometry which has tours T^d . Before imposing these conditions, let us compare the gravity couplings that the Z_2 -symmetry produces, with the couplings in the Euler character I_2 .

3.2.1 Comparing with the Euler character I_2

In this subsection we compare the couplings in the classical effective action that are consistent with the T-duality, with the couplings in the Chern–Simons form Q_2 in the Euler character I_2 . When the dilaton and B -field are zero, the couplings that are found by the T-duality constraint are the following:

$$\begin{aligned}
 &+2b_{19} \left(K^\gamma_\gamma n^\alpha n^\beta R_{\alpha\beta} - 12b_1 K^\alpha_\alpha \right. \\
 &+ b_{11} K^{\gamma\delta} n^\alpha n^\beta R_{\alpha\gamma\beta\delta} \\
 &\left. + b_{38} n^\alpha n^\beta n^\gamma n^\delta \nabla_\delta \nabla_\gamma K_{\alpha\beta} \right]
 \end{aligned}$$

$$\begin{aligned}
 1 + \partial S_1 = & -\frac{48a_1}{\kappa^2} \int d^D x \sqrt{-G} (R^{\mu\nu\alpha\beta} R_{\mu\nu\alpha\beta} \\
 & - 4R^{\mu\nu} R_{\mu\nu} + R^2) \\
 & -\frac{2}{\kappa^2} \int d^{D-1} x \sqrt{|g|} \left[b_{11} K_{\alpha\gamma} K^{\alpha\beta} K_{\beta\gamma} \right. \\
 & + \left(24b_1 - \frac{1}{2}b_{17} \right) K^\alpha_\alpha K_{\beta\gamma} K^{\beta\gamma} \\
 & + \left(-8b_1 - \frac{1}{6}b_{18} \right) K^\alpha_\alpha K^\beta_\beta K^\gamma_\gamma \\
 & + 12(-16a_1 + b_{12}) K^{\alpha\beta} R_{\alpha\beta} \\
 & \left. + \left(96a_1 + 24b_1 - 12b_{12} - \frac{1}{2}b_{17} \right) \right]
 \end{aligned}$$

R

(56)

where the couplings are all independent. One may use various identities to rewrite the boundary coupling in different form, however, the number of boundary couplings remain the same. For example, one may use the following identity:

$$n^\alpha n^\beta n^\gamma n^\delta \nabla_\delta \nabla_\gamma K_{\alpha\beta} = -2K_{\alpha\gamma} K_{\alpha\beta} K_{\beta\gamma} + n^\alpha n^\beta \nabla_\gamma \nabla_\gamma K_{\alpha\beta}$$

(57)

which can be verified by writing both sides in terms of function f , to write the term with the second derivative of extrinsic curvature in (56) in terms of the Laplacian of the extrinsic curvature. This changes the coefficient of $K^\alpha_\alpha K^{\beta\gamma} K_{\beta\gamma}$, however, the number of the boundary couplings remain the same.

On the other hand, it has been shown in [27] that if one extends the couplings in the four-dimensional Euler density I_2 to arbitrary dimension, the couplings satisfy the least action principle with the boundary condition that only metric is arbitrary on the boundary. This extension is

$$I_2 = \int d^D x \sqrt{-G} R_{GB}^2 + \int d^{D-1} x \sqrt{|g|} Q_2 \tag{58}$$

where R_{GB}^2 is the Gauss–Bonnet gravity and Q_2 is the Chern–Simons form [27]

$$\begin{aligned}
 2 = 4 \left[K^\mu_\mu \bar{R} - 2K^{\mu\nu} \bar{R}_{\mu\nu} + \frac{1}{3} (3K^\alpha_\alpha K_{\mu\nu} K^{\mu\nu} \right. \\
 \left. - K^\mu_\mu K^\nu_\nu K^\alpha_\alpha - 2K_{\mu\nu} K_{\nu\alpha} K^{\alpha\mu}) \right] Q
 \end{aligned} \tag{59}$$

where $R^{\sim}_{\mu\nu}$ and R^{\sim} are curvatures that are constructed from the induced metric (9). Using the following Gauss–Codazzi relations:

$$R^{\sim\alpha\beta} = P_{\alpha\mu}P_{\beta\nu}R_{\mu\nu} - n_{\mu\nu}R_{\alpha\mu\beta\nu} - K_{\alpha\mu}K_{\beta\mu} + K_{\alpha\beta}K_{\mu\mu}$$

$$R^{\sim} = R - 2n_{\mu\nu}R_{\mu\nu} - K_{\mu\nu}K_{\mu\nu} + K_{\mu\mu}K_{\nu\nu} \quad (60)$$

and the identity $n^{\mu}K_{\mu\nu} = 0$, one can rewrite Q_2 in terms of the spacetime curvatures, i.e.,

$$Q_2 = 4 \left[K^{\mu}_{\mu} R - 2K^{\mu\nu} R_{\mu\nu} - 2K_{\alpha}^{\alpha} n^{\mu} n^{\nu} R_{\mu\nu} \right. \\ \left. + 2K^{\mu\nu} n^{\alpha} n^{\beta} R_{\alpha\mu\beta\nu} \right. \\ \left. - \frac{1}{3} (6K^{\alpha}_{\alpha} K_{\mu\nu} K^{\mu\nu} - 2K^{\mu}_{\mu} K^{\nu}_{\nu} K^{\alpha}_{\alpha} \right. \\ \left. - 4K_{\mu}^{\nu} K_{\nu\alpha} K^{\alpha\mu} \right] \quad (61)$$

The bulk couplings in I_2 are the same as the bulk couplings in (56), however, the number of boundary couplings in (56) is more than the number of boundary couplings in I_2 . If one sets b_{38} to zero in (56), then the number of couplings and the structure of couplings become the same as those in I_2 , however, for no values of the boundary parameters $b_1, b_{11}, b_{12}, b_{17}, b_{18}, b_{19}$ the two sets of the couplings become identical, e.g., the ratio of the last terms in the first and second lines above is 3/2 whereas this ratio in (56) is one. Hence, the boundary couplings that the T-duality dictates can not be *exactly* the same as the boundary couplings in I_2 for any values of the boundary parameters. This may indicate that the assumption that only metric is arbitrary on the boundary in the 4-derivative couplings is not consistent with the classical effective action of string theory. However, such assumption may be valid for the higher loop effective actions for which there is no T-duality symmetry.

4 Constraint from the least action principle

We have seen that when the background geometry has a circle, requiring the bulk and boundary actions at order α to have the Z_2 -symmetry, one finds the bulk action (45) and the boundary action (55). Using the background independent assumption, these actions should be the effective actions at

order α for any arbitrary background which has boundary, up to field redefinitions. To find the bulk equations of motion one needs to extremize these actions, i.e., $\delta(\mathbf{S}_1 + \partial\mathbf{S}_1) = 0$, by using some assumption for the massless fields on the boundary. At the two-derivative order, the assumption that the massless fields should be arbitrary on the boundary is good enough to produce the equations of motion from extremizing the leading order actions, i.e., $\delta(\mathbf{S}_0 + \partial\mathbf{S}_0) = 0$. At the four-derivative order, however, the assumption that only the massless fields are arbitrary requires the boundary couplings which are not consistent with the T-duality symmetry. If one insists on that assumption, then the 4-derivative action would not be the classical effective action of the string theory in which the T-duality is a symmetry. Hence, it seems to study the classical equations of motion in the string theory at order α , one should assume that not only the massless fields but also their first derivatives should be arbitrary on the boundary.

Since the bulk action (45) has at most two-derivative terms, e.g., R or $\nabla\nabla\Phi$, when one extremizes the bulk action, one would find the variation δ as well as the variations $\nabla(\delta\Psi)$ and $\nabla\nabla(\delta)$ where Ψ represents the massless fields. After using the Stokes's theorem, the variations $\nabla(\delta)$ and $\nabla\nabla(\delta)$ produce the variations $\delta\Psi$ and $\nabla(\delta)$ on the boundary. The assumption that Ψ and $\nabla\Psi$ are arbitrary on the boundary, means their variations are zero on the boundary, i.e., $\delta\Psi = \nabla(\delta\Psi) = 0$. Hence, the bulk action satisfies $\delta\mathbf{S}_1 = (\dots)\delta\Psi = 0$ with no constraint on the bulk couplings. The boundary action (55), however, has two- and three-derivative terms, e.g., R or $\nabla\nabla K$. When one extremizes the boundary action, one would find variations $\nabla\nabla(\delta)$ and $\nabla\nabla\nabla(\delta)$ on the boundary which are not zero in general. The parameters in the boundary action should be such that the coefficients of these variations become zero up to some total derivative terms on the boundary which are zero according to the Stokes's theorem.

The variation of the boundary action (55) against the metric variation produces the following non-zero terms in the local frame:

$$24(8a_1 + b_1)\partial^{\alpha}\Phi f^1{}^{\beta} P^{\gamma\delta} \partial_{\alpha}\partial_{\beta}\delta G_{\gamma\delta} \\ + (96a_1 - \frac{1}{2}b_{11} - 6b_{12})f^1{}^{\alpha} f^1{}^{\beta} f^2{}^{\gamma\delta} P_{\gamma}{}^{\epsilon} P_{\delta}{}^{\epsilon} \partial_{\alpha}\partial_{\beta}\delta G_{\epsilon\epsilon} \\ + (-48a_1 + 6b_{12} + \frac{1}{4}b_{17} - b_{19}) \\ \times f^1{}^{\alpha} f^1{}^{\beta} f^2{}^{\gamma\delta} P_{\gamma\delta} P^{\epsilon\epsilon} \partial_{\alpha}\partial_{\beta}\delta G_{\epsilon\epsilon} \\ + (-96a_1 - 24b_1 - 12b_{12} - \frac{1}{2}b_{17} + 2b_{19})\partial^{\alpha}\Phi \\ \times f^1{}_{\alpha} f^1{}^{\beta} f^1{}^{\gamma} P^{\delta\epsilon} \partial_{\beta}\partial_{\gamma}\delta G_{\delta\epsilon} \quad (62)$$

where $f^1 = \partial^\alpha f$, $f^2 = \partial^\alpha \partial^\beta f$. We have used the assumption that the variation of metric and its first derivative, and their tangent derivatives are zero, i.e., $\delta G_{\alpha\beta} = \partial_\mu \delta G_{\alpha\beta} = 0$ and $P^{\mu\nu} \partial_\mu \partial_\nu \delta G_{\alpha\beta} = 0$. One is free to add arbitrary total derivative terms, i.e., (16) in which the antisymmetric tensor $\mathcal{F}_1^{\alpha\beta}$ contains the variation of metric. Using the Stokes's theorem, the total derivative terms on the boundary become zero. Then, up to some total derivative terms, the resulting equations are zero for the following relation:

$$b_{19} = -48a_1 + 6b_{12} + b_{17}/4 \tag{63}$$

Note that if one requires that the first derivative of the variation of metric to be non-zero on the boundary, i.e., $\partial_\mu \delta G_{\alpha\beta} = 0$, then one would find many other terms in (62) that become zero for the incorrect value of $a_1 = 0$.

Inserting the relation (63) into (55), one finds the variation of (55) against the dilaton becomes zero up to some total derivative terms, and the B -field variation produces the following relation:

$$b_{12} = 16a_1 - b_{11}/12, \tag{64}$$

and some total derivative terms. Therefore, for the relations (63) and (64) the bulk and boundary actions satisfy the stationary condition $\delta(\mathbf{S}_1 + \partial\mathbf{S}_1) = (\dots)\delta\Psi = 0$ when the variation of massless fields and their first derivative on the boundary are zero.³ Since the variation of fields in the bulk are non-zero, this gives the appropriate equations of motion in which we are not interested.

Inserting the relations (63) and (64) into the boundary action (55), one finds the boundary action to be

$$\begin{aligned} \partial\mathcal{L}_1 = & b_{11} \left[K_\alpha^\gamma K^{\alpha\beta} K_{\beta\gamma} + \frac{1}{4} H_\alpha^{\delta\epsilon} H_{\beta\delta\epsilon} K^\gamma{}_\gamma \right. \\ & - K^{\alpha\beta} R_{\alpha\beta} + K^{\gamma\delta} n^\alpha n^\beta R_{\alpha\gamma\beta\delta} \\ & - \frac{1}{12} H^{\beta\gamma\delta} n^\alpha \nabla_\alpha H_{\beta\gamma\delta} - \frac{1}{2} H_\beta^{\delta\epsilon} H_{\gamma\delta\epsilon} n^\alpha n^\beta n^\gamma \nabla_\alpha \Phi \\ & \left. - 2K^{\alpha\beta} \nabla_\beta \nabla_\alpha \Phi + \frac{1}{4} H_\alpha^{\delta\epsilon} n^\alpha n^\beta n^\gamma \nabla_\gamma H_{\beta\delta\epsilon} \right] \\ & + b_1 \left[H_{\beta\gamma\delta} H^{\beta\gamma\delta} K^\alpha{}_\alpha + 24K^\alpha{}_\alpha K_{\beta\gamma} K^{\beta\gamma} \right. \\ & \left. - 8K_{\alpha\alpha} K_{\beta\beta} K_{\gamma\gamma} + 24K_{\gamma\gamma} n_\alpha n_\beta R_{\alpha\beta} \right] \end{aligned}$$

$$\begin{aligned} & - 2H_{\beta\gamma\delta} H^{\beta\gamma\delta} n^\alpha \nabla_\alpha \Phi + 24n^\alpha R \nabla_\alpha \Phi \\ & - 48K^\beta{}_\beta \nabla_\alpha \Phi \nabla^\alpha \Phi \\ & 12K_\alpha R \\ & + 96K^\gamma{}_\gamma n^\alpha n^\beta \nabla_\alpha \Phi \nabla_\beta \Phi \\ & - 48K^{\alpha\beta} \nabla_\beta \nabla_\alpha \Phi - 48n^\alpha R_{\alpha\beta} \nabla^\beta \Phi \\ & + 96n^\alpha \nabla_\alpha \Phi \nabla_\beta \Phi \nabla^\beta \Phi \\ & - 96n^\alpha \nabla_\beta \nabla_\alpha \Phi \nabla^\beta \Phi \\ & - 128n_\alpha n_\beta n_\gamma \nabla_\alpha \Phi \nabla_\beta \Phi \nabla_\gamma \Phi \\ & + 96n^\alpha n^\beta n^\gamma \nabla_\alpha \Phi \nabla_\gamma \nabla_\beta \Phi \Big] \\ & + a_1 \left[24H_\alpha^{\gamma\delta} H_{\beta\gamma\delta} K^{\alpha\beta} - 24H_\alpha^{\delta\epsilon} H_{\beta\delta\epsilon} K^\gamma{}_\gamma n^\alpha n^\beta \right. \\ & + 16H^{\beta\gamma\delta} n^\alpha \nabla_\alpha H_{\beta\gamma\delta} \\ & + 48H_\beta^{\delta\epsilon} H_{\gamma\delta\epsilon} n_\alpha n_\beta n^\gamma \nabla_\alpha \Phi + 192n^\alpha R \nabla_\alpha \Phi \\ & + 384K^\gamma{}_\gamma n^\alpha n^\beta \nabla_\alpha \Phi \nabla_\beta \Phi - 384n^\alpha R_{\alpha\beta} \nabla^\beta \Phi \\ & - 384K_{\alpha\beta} \nabla^\alpha \Phi \nabla^\beta \Phi + 384n^\alpha \nabla_\alpha \Phi \nabla_\beta \Phi \nabla^\beta \Phi \\ & - 768n^\alpha \nabla_\beta \nabla_\alpha \Phi \nabla^\beta \Phi \\ & - 48H_\alpha n^\beta n^\gamma \nabla_\gamma H_{\beta\delta\epsilon} \\ & \left. - 768n_\alpha n_\beta n_\gamma \nabla_\alpha \Phi \nabla_\beta \Phi \nabla_\gamma \Phi \right. \\ & \left. + 768n^\alpha n^\beta n^\gamma \nabla_\alpha \Phi \nabla_\gamma \nabla_\beta \Phi \right] \\ & - \frac{b_{17}}{2} \left[K^\alpha{}_\alpha K_{\beta\gamma} K^{\beta\gamma} + \frac{1}{4} H_\alpha^{\delta\epsilon} H_{\beta\delta\epsilon} K^\gamma{}_\gamma n^\alpha n^\beta \right. \\ & \left. - K_{\beta\gamma} K^{\beta\gamma} n^\alpha \nabla_\alpha \Phi - \frac{1}{2} H_\beta^{\delta\epsilon} H_{\gamma\delta\epsilon} n^\alpha n^\beta n^\gamma \nabla_\alpha \Phi \right] \\ & - \frac{b_{18}}{6} \left[K^\alpha{}_\alpha - 2n^\alpha \nabla_\alpha \Phi \right]^3 \\ & + b_{38} n^\alpha n^\beta n^\gamma n^\delta \nabla_\delta \nabla_\gamma K_{\alpha\beta} \tag{65} \end{aligned}$$

The boundary multiplets with the background independent parameters $b_1, b_{11}, b_{17}, b_{18}$ and b_{38} are each invariant under the Z_2 -symmetry and satisfy the least action. The boundary multiplet with parameter a_1 also satisfies the least action principle, however, its combination with the bulk multiplet (45) satisfies the Z_2 -symmetry. In the next section we study another constraint on the background independent parameters by considering the geometry which has the torsion T^d .

³ If one does not use the total derivative terms (16) in the metric variation of the boundary couplings (62), then one would find the constraint (63), (64), and another constraint $b_1 = -8a_1$.

5 Constraint from zero cosmological boundary action

In this section we show that the cosmological reduction of the boundary term at the leading order produces zero boundary action. We then extend this to the cosmological boundary action at order α to further constrain the boundary parameters in (65).

When fields depend only on time, using the gauge symmetries it is possible to write the metric, B -field and dilaton as

$$\begin{aligned} G_{\mu\nu} &= \begin{pmatrix} -n^2(t) & 0 \\ 0 & G_{ij}(t) \end{pmatrix}, \\ B_{\mu\nu} &= \begin{pmatrix} 0 & 0 \\ 0 & B_{ij}(t) \end{pmatrix}, \\ 2\Phi &= \phi + \frac{1}{2} \log \det(G_{ij}) \end{aligned} \quad (66)$$

where the lapse function $n(t)$ can also be fixed to $n = 1$. The cosmological reduction of the bulk action in (6) then becomes

$$\begin{aligned} \mathbf{S}_0^c &= -\frac{2}{\kappa^2} \int dt e^{-\phi} \left[\frac{1}{4} B_{ij} B^{ij} \right. \\ &\quad \left. - \frac{3}{4} G^i{}_{ij} G^j{}_{ij} - G^{ij} G^i{}_{ij} \phi' - \phi'^2 + G^{ij} G^{ij} \right] \end{aligned} \quad (67)$$

where $G^i{}_{ij} \equiv G^{ik} G^{jl} G^k{}_{kl}$. Up to a total derivative term the above action can be written in $O(d,d)$ -invariant form. In fact, using the following total derivative term:

$$\begin{aligned} & - \left[e^{-\phi} \frac{d}{dt} G^i{}_{ij} \right. \\ & \left. \int dt e^{-\phi} \left[- \frac{1}{4} G^{ij} G^i{}_{ij} \phi' - G^i{}_{ij} G^j{}_{ij} + G^{ij} G^{ij} \right] \right] \end{aligned} \quad (68)$$

one can write \mathbf{S}_0^c as

$$\begin{aligned} \mathbf{S}_0^c &= -\frac{2}{\kappa^2} \int dt e^{-\phi} \left[\frac{1}{4} B_{ij} B^{ij} + \frac{1}{4} G^i{}_{ij} G^j{}_{ij} - \phi'^2 \right] \\ & - \int dt \left[e^{-\phi} \frac{d}{dt} G^i{}_{ij} - \kappa^2 \frac{d}{dt} G^{ij} \right] \end{aligned} \quad (69)$$

Since there is boundary, the total derivative term can not be ignored. It can be transferred to the boundary by using the Stokes's theorem.

On the other hand, the cosmological boundary is specified by $x^i = \sigma^i$, and $x^0 = t$ is independent of σ^i . Hence, the cosmological reduction of $\sqrt{|g|} e^{2\Phi} = e^{-\phi}$. The unit vector to the boundary is fixed, i.e., $n^i = 0$, and the cosmological reduction of the trace of the extrinsic curvature becomes

$$K = \frac{1}{2} G^i{}_{ij} G^j{}_{ij} \quad (70)$$

Therefore, the reduction of the boundary term in (6) is exactly cancelled with the total derivative term in the bulk action, i.e.,

$$\begin{aligned} \mathbf{S}_0^c &= -\frac{2}{\kappa^2} \int dt e^{-\phi} \left[\frac{1}{4} \text{Tr} W^2 - \frac{1}{4} \text{Tr} Y^2 - \phi'^2 \right] \\ \partial \mathbf{S}_0^c &= 0 \end{aligned} \quad (71)$$

where $W = G^{-1} G^i{}_{ij}$, $Y = G^{-1} B^i{}_{ij}$. The bulk action is invariant under $O(d,d)$ [5,6]. Note that the reduction of the extrinsic curvature (70) can not be written in $O(d,d)$ invariant form. So it was necessary that this term was cancelled with the total derivative term in the bulk action. In other words, there is no way to write the boundary action in $O(d,d)$ invariant form unless it is zero.

At the higher order of α the bulk and boundary actions should be invariant under $O(d,d)$. We will see the couplings that are found by the Z_2 -symmetry satisfy the $O(d,d)$ symmetry with no further constraint on the boundary parameters. However, since the cosmological reduction of the leading order boundary action is zero we speculate that the boundary action at all higher orders of α to be zero too, i.e., (7). This can be a consistency check and a constraint for the boundary couplings at higher orders of α , e.g., the cosmological reduction of the boundary couplings (65) should satisfy the $O(d,d)$ symmetry and then the constraint (7) fixes the coefficients of the $O(d,d)$ invariant terms to be zero.

Using a one-dimensional field-redefinition for which the lapse function remains invariant,⁴ it has been shown in [8] that the cosmological reduction of the bulk action (45) can

⁴ Note that the lapse function in the cosmological reduction corresponds to the base space metric g^{-ab} in the circular reduction. The invariance of the lapse function is then consistent with the invariance of the base

space metric under the Z_2 -transformations that we have considered in this paper.

be written explicitly in $O(d,d)$ -invariant form when there is no boundary, i.e.,

$$S_1^c = -\frac{2}{\kappa^2} 24a_1 \int dt e^{-\phi} \left[\frac{1}{8} \text{Tr}(Y^4) + \frac{1}{8} \text{Tr}(W^4) + \frac{1}{4} \text{Tr}_{(YWYW)} - \frac{1}{2} \text{Tr}(Y^2 W^2) - \frac{1}{16} [\text{Tr}(Y^2) - \text{Tr}(W^2)]^2 - \frac{1}{2} [\text{Tr}(Y^2) - \text{Tr}(W^2)] \dot{\phi}^2 - \frac{1}{3} \dot{\phi}^4 \right] \tag{72}$$

In this calculation the terms which could not be written in $O(d,d)$ -invariant form are removable by field redefinition and total derivative terms. The total derivative terms which are needed to write the cosmological reduction of the bulk couplings in the above $O(d,d)$ -invariant form, are the following:

$$24a_1 \left[\frac{e^{-\phi}}{dt} \frac{\text{Tr}(WY^2)}{dt} - \frac{e^{-\phi}}{2} \frac{\dot{\phi}^3}{dt} + \frac{e^{-\phi}}{4} \frac{\dot{\phi}^2 \text{Tr}(W)}{dt} + \frac{1}{2} e^{-\phi} \dot{\phi} (\text{Tr} W)^2 - \frac{1}{2} e^{-\phi} \text{Tr}(W)^2 \text{Tr}(W) + \frac{1}{4} e^{-\phi} (\text{Tr} W)^3 \right] \tag{73}$$

These total derivative terms have been ignored in [8] because it has been assumed that the spacetime has no boundary. In the presence of boundary, the above total derivative terms produce the following boundary terms:

$$24a_1 n e^{-\phi} \left[\text{Tr}(WY^2) + \frac{2}{3} \dot{\phi}^3 + \dot{\phi}^2 \text{Tr}(W) + \frac{1}{2} \dot{\phi} (\text{Tr} W)^2 - \frac{1}{2} \text{Tr}(W)^2 \text{Tr}(W) + \frac{1}{4} (\text{Tr} W)^3 \right] \tag{74}$$

which should be taken into account when studying the $O(d,d)$ -symmetry of the boundary action (65). The boundary of the cosmological reduction is spacelike. The one-dimensional reduction that its boundary is timelike is the same as above in which $n = -1$.

On the other hand, using the reductions (66), one finds the cosmological reduction of the boundary action (65) for the following relation between b_1, b_{11} :

$$b_{11} = -24b_1$$

to be

$$24a_1 n e^{-\phi} \left[-\text{Tr}(WY^2) - \frac{2}{3} \dot{\phi}^3 - \dot{\phi}^2 \text{Tr}(W) - \frac{1}{2} \dot{\phi} (\text{Tr} W)^2 + \frac{1}{2} \text{Tr}(W)^2 \text{Tr}(W) - \frac{1}{4} (\text{Tr} W)^3 \right] + n e^{-\phi} \left[(24a_1 + 3b_1 + \frac{b_{17}}{8}) \dot{\phi} (\text{Tr} W^2 - \text{Tr} Y^2) + (32a_1 + 4b_1 - \frac{1}{6} b_{18}) \dot{\phi}^3 \right] \tag{75}$$

Note that the cosmological reduction of the coupling in (65) with coefficient b_{38} is zero. While the terms in the second line above are invariant under the $O(d,d)$ transformations, the terms in the first line are not. However, adding the residual total derivative terms from the bulk action, i.e., (74), one finds the boundary terms in the first line above are cancelled. Note that if one changes the coefficient of the boundary coupling $K_\alpha^\nu K^{\alpha\beta} K_{\beta\nu}$ in (65), then there would be the term $\text{Tr}(W)^3$ in the first bracket above which is not cancelled with the total derivative terms and is not invariant under $O(d,d)$ transformation.

It means the cosmological reduction of the Euler character is not consistent with the $O(d,d)$ symmetry.

Therefore, up to a field redefinition, the cosmological reduction of the bulk and boundary couplings are given by the $O(d,d)$ invariant bulk action (73) and the following $O(d,d)$ invariant boundary action:

$$\partial S_1^c = -\frac{2n}{\kappa^2} e^{-\phi} \left[(-24a_1 - 3b_1 - \frac{b_{17}}{8}) \dot{\phi} (\text{Tr} W^2 - \text{Tr} Y^2) + (32a_1 + 4b_1 - \frac{b_{18}}{6}) \dot{\phi}^3 \right] \tag{76}$$

Note that since the cosmological boundary action at the leading order is zero, i.e., (71), the field redefinition has no effect on the above cosmological boundary action. Hence, requiring the effective actions that are found by the Z_2 -symmetry and by the least action principle, to be invariant under the $O(d,d)$ -transformations produces no further constraint on the boundary couplings. However, requiring the constraint (7), one finds the following two relations between the parameters.

$$\begin{aligned}
 S_1 + \partial S_1|_{\Phi=B=0} &= -\frac{48a_1}{\kappa^2} \left[\int d^D x \sqrt{-G_{\text{RGB2}}} \right. \\
 &\quad \left. + \int d^{D-1} x \sqrt{|g|} \left(Q_2 + \frac{4}{3} n^\alpha n^\beta \nabla_\gamma \nabla^\gamma K_{\alpha\beta} \right) \right] \quad (80)
 \end{aligned}$$

While the Euler character I_2 is not consistent with the symmetries of the classical effective actions, i.e., it does not satisfy the $O(1,1)$ symmetry when the geometry has one circle, nor with $O(d,d)$ symmetry when the geometry has the tours T^d , the above couplings are consistent with the $O(1,1)$ and $O(d,d)$ symmetries.

The effective actions that we have found for the relations (79) are the following:

$$\begin{aligned}
 I_1 &= -\frac{48a_1}{\kappa^2} \int_M d^D x \sqrt{-G} e^{-2\Phi} \\
 &\quad \times \left[R_{GB}^2 + \frac{1}{24} H_\alpha^{\delta\epsilon} H^{\alpha\beta\gamma} H_{\beta\delta\epsilon} H_{\gamma\epsilon\delta} \right. \\
 &\quad - \frac{1}{8} H_{\alpha\beta}^{\delta\epsilon} H^{\alpha\beta\gamma} H_{\gamma\epsilon\delta} H_{\delta\epsilon\gamma} \\
 &\quad + \frac{1}{144} H_{\alpha\beta\gamma} H^{\alpha\beta\gamma} H_{\delta\epsilon\gamma} H^{\delta\epsilon\gamma} + H_\alpha^{\gamma\delta} H_{\beta\gamma\delta} R^{\alpha\beta} \\
 &\quad - \frac{1}{6} H_{\alpha\beta\gamma} H^{\alpha\beta\gamma} R - \frac{1}{2} H_\alpha^{\delta\epsilon} H^{\alpha\beta\gamma} R_{\beta\gamma\delta\epsilon} \\
 &\quad - \frac{2}{3} H_{\beta\gamma\delta} H^{\beta\gamma\delta} \nabla_\alpha \nabla^\alpha \Phi + \frac{2}{3} H_{\beta\gamma\delta} H^{\beta\gamma\delta} \nabla_\alpha \Phi \nabla^\alpha \Phi \\
 &\quad + 8R \nabla_\alpha \Phi \nabla^\alpha \Phi + 16 \nabla_\alpha \Phi \nabla^\alpha \Phi \nabla_\beta \nabla^\beta \Phi \\
 &\quad - 16 R_{\alpha\beta} \nabla^\alpha \Phi \nabla^\beta \Phi - 16 \nabla_\alpha \Phi \nabla^\alpha \Phi \nabla_\beta \nabla^\beta \Phi \\
 &\quad \left. + 2H_\alpha^{\gamma\delta} H_{\beta\gamma\delta} \nabla^\beta \nabla^\alpha \Phi \right] \quad (81) \\
 &= -\frac{a_1}{2} \int_M d^D x \sqrt{|g|} e^{-2\Phi} \quad 48
 \end{aligned}$$

$$\begin{aligned}
 \partial S_1 &= \int_{\partial M} d^D x \sqrt{|g|} \kappa \\
 &\quad \times \left[Q_2 + \frac{4}{3} n^\alpha n^\beta \nabla_\gamma \nabla^\gamma K_{\alpha\beta} \right. \\
 &\quad + \frac{2}{3} H_{\beta\gamma\delta} H^{\beta\gamma\delta} n^\alpha \nabla_\alpha \Phi \\
 &\quad - 2H_\beta^{\delta\epsilon} H_{\gamma\delta\epsilon} n^\alpha n^\beta \nabla_\alpha \Phi \\
 &\quad - \frac{1}{3} H_{\beta\gamma\delta} H^{\beta\gamma\delta} K^\alpha_\alpha + H_\alpha^{\gamma\delta} H_{\beta\gamma\delta} K^{\alpha\beta} \\
 &\quad + H_\alpha^{\delta\epsilon} H_{\beta\delta\epsilon} K^\gamma_\gamma n^\alpha n^\beta \\
 &\quad - 16K^\gamma_\gamma n^\alpha n^\beta \nabla_\alpha \Phi \nabla_\beta \Phi + 16K^\beta_\beta \nabla_\alpha \Phi \nabla^\alpha \Phi \\
 &\quad - 16K_{\alpha\beta} \nabla^\alpha \Phi \nabla^\beta \Phi \\
 &\quad \left. - 16n^\alpha \nabla_\alpha \Phi \nabla_\beta \Phi \nabla^\beta \Phi + \frac{32}{3} n^\alpha n^\beta n^\gamma \nabla_\alpha \Phi \nabla_\beta \Phi \nabla_\gamma \Phi \right] \quad (82)
 \end{aligned}$$

where R_{GB}^2 is the Gauss–Bonnet bulk couplings and Q_2 is the Chern–Simons boundary couplings (61). The bulk couplings for $a_1 = 1/96$ is the effective action of the bosonic string theory which has been found in [8].

We have imposed the relation (44) to have standard propagators for the B-field. This relation can be also found by the $O(d,d)$ symmetry. We have seen that the cosmological reduction of the bulk couplings are invariant under the $O(d,d)$ transformation up to some total derivative terms which are not invariant. These anomalous terms are exactly cancelled with the anomalous terms in the cosmological reduction of the boundary couplings. If one does not use the constraint (44), then the two set of anomalous terms would cancel each other only under the condition (44). We have performed this calculation explicitly.

In the cosmological study, we have used the scheme that the cosmological action has the first derivative of dilaton, i.e., (72), and the boundary action has no term with first derivative of dilaton, i.e., (7). On the other hand, it has been shown in [31,32] that if one uses various one-dimensional field redefinitions and uses integration by part, then the cosmological reduction of the bulk action at order α and higher can be written in a scheme in which the bulk action has only the first derivative of the generalized metric, i.e., no coupling involves the first derivative of dilaton. In the presence of boundary, the total derivative terms appear in the boundary by using the Stokes’s theorem. Hence, if one uses the scheme in which the derivative of dilaton does not appear in the bulk action, then in that scheme the boundary action may have the first derivative of the dilaton, i.e., the cosmological boundary action may not be zero in that scheme.

In imposing the $O(1,1)$ symmetry, we have assumed the unit normal vector of the boundary in the base space, n^a and its length remain invariant under the T-duality transformations. This forces us to work with the most general gauge invariant bulk action (12) which has 20 parameters, i.e., we did not use the higher derivative field redefinitions to work with the independent bulk couplings. If one uses the most general field redefinitions, then the bulk action in the minimal scheme has only 8 independent couplings. The T-duality fixes these parameters up to an overall factor [11], i.e.,

$$\begin{aligned}
 S^1 &= \frac{-2c_1}{\kappa^2} \int_M d^D x e^{-2\Phi} \sqrt{-G} \\
 &\quad \times \left(R_{\alpha\beta\gamma\delta} R^{\alpha\beta\gamma\delta} - \frac{1}{2} H_\alpha^{\delta\epsilon} H^{\alpha\beta\gamma} R_{\beta\gamma\delta\epsilon} \right)
 \end{aligned}$$

$$+\frac{1}{24}H_{\epsilon\delta\zeta}H^{\epsilon\beta}H^{\delta\gamma}H^{\zeta\alpha}-\frac{1}{8}H_{\alpha\beta}{}^{\delta}H^{\alpha\beta\gamma}H_{\gamma}{}^{\epsilon\zeta}H_{\delta\epsilon\zeta}) \quad (83)$$

where c_1 is the overall factor. For $c_1 = 1/4$, the above action is the effective action of the bosonic string theory at order α which has been found in [12] by the S-matrix method. The above action and the action (81) are related to each other by a particular field redefinition [8]. In the presence of boundary, however, one may not be able to use the most general field redefinitions because they change the values of the massless fields and their derivatives on the boundary which may not be consistent with the least action principle. It would be interesting to find the appropriate field redefinitions in the presence of the boundary to find the corresponding independent gauge invariant couplings and then impose the constraints that we have studied in this paper, to find independent bulk and boundary couplings at order α . It would be also interesting to extend the calculation in this paper to find the boundary couplings at order α^2, α^3 . The corresponding bulk actions in the minimal scheme have been found in [13, 16, 17].

Data Availability Statement This manuscript has no associated data or the data will not be deposited. [Authors' comment: The results in this paper are obtained analytically, hence, it does not use any data.]

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. Funded by SCOAP³.

References

- J.M. Maldacena, Int. J. Theor. Phys. **38**, 1113 (1999). <https://doi.org/10.1023/A:1026654312961>
- J.M. Maldacena, Adv. Theor. Math. Phys. **2**, 231 (1998). <https://doi.org/10.4310/ATMP.1998.v2.n2.a1> arXiv:hep-th/9711200
- J.W. York Jr., Phys. Rev. Lett. **28**, 1082 (1972). <https://doi.org/10.1103/PhysRevLett.28.1082>
- G.W. Gibbons, S.W. Hawking, Phys. Rev. D **15**, 2752 (1977). <https://doi.org/10.1103/PhysRevD.15.2752>
- G. Veneziano, Phys. Lett. B **265**, 287–294 (1991). [https://doi.org/10.1016/0370-2693\(91\)90055-U](https://doi.org/10.1016/0370-2693(91)90055-U)
- K.A. Meissner, G. Veneziano, Phys. Lett. B **267**, 33–36 (1991). [https://doi.org/10.1016/0370-2693\(91\)90520-Z](https://doi.org/10.1016/0370-2693(91)90520-Z)
- J. Maharana, J.H. Schwarz, Nucl. Phys. B **390**, 3–32 (1993). [https://doi.org/10.1016/0550-3213\(93\)90387-5](https://doi.org/10.1016/0550-3213(93)90387-5) arXiv:hep-th/9207016
- K.A. Meissner, Phys. Lett. B **392**, 298–304 (1997). [https://doi.org/10.1016/S0370-2693\(96\)01556-0](https://doi.org/10.1016/S0370-2693(96)01556-0) arXiv:hep-th/9610131
- A. Sen, Phys. Lett. B **271**, 295–300 (1991). [https://doi.org/10.1016/0370-2693\(91\)90090-D](https://doi.org/10.1016/0370-2693(91)90090-D)
- O. Hohm, A. Sen, B. Zwiebach, JHEP **02**, 079 (2015). [https://doi.org/10.1007/JHEP02\(2015\)079](https://doi.org/10.1007/JHEP02(2015)079) arXiv:1411.5696 [hep-th]
- M.R. Garousi, Phys. Rev. D **99**(12), 126005 (2019). <https://doi.org/10.1103/PhysRevD.99.126005> arXiv:1904.11282 [hep-th]
- R.R. Metsaev, A.A. Tseytlin, Nucl. Phys. B **293**, 385 (1987). [https://doi.org/10.1016/0550-3213\(87\)90077-0](https://doi.org/10.1016/0550-3213(87)90077-0)
- M.R. Garousi, Eur. Phys. J. C **79**(10), 827 (2019). <https://doi.org/10.1140/epjc/s10052-019-7357-4> arXiv:1907.06500 [hep-th]
- H. Razaghian, M.R. Garousi, Phys. Rev. D **97**, 106013 (2018). <https://doi.org/10.1103/PhysRevD.97.106013> arXiv:1801.06834 [hep-th]
- M.R. Garousi, Eur. Phys. J. C **80**, 1086 (2020). <https://doi.org/10.1140/epjc/s10052-020-08662-9> arXiv:2006.09193 [hep-th]
- M.R. Garousi, JHEP **02**, 157 (2021). [https://doi.org/10.1007/JHEP02\(2021\)157](https://doi.org/10.1007/JHEP02(2021)157) arXiv:2011.02753 [hep-th]
- M.R. Garousi, Nucl. Phys. B **971**, 115510 (2021). <https://doi.org/10.1016/j.nuclphysb.2021.115510> arXiv:2012.15091 [hep-th]
- M.R. Garousi, Phys. Rev. D **104**(6), 066013 (2021). <https://doi.org/10.1103/PhysRevD.104.066013> arXiv:2105.07598 [hep-th]
- M.R. Garousi, Eur. Phys. J. C **81**(8), 711 (2021). <https://doi.org/10.1140/epjc/s10052-021-09518-6> arXiv:2107.04796 [hep-th]
- M.R. Garousi, Phys. Rep. **702**, 1 (2017). <https://doi.org/10.1016/j.physrep.2017.07.009> arXiv:1702.00191 [hep-th]
- M. Mashhadi, M.R. Garousi, JHEP **06**, 171 (2020). [https://doi.org/10.1007/JHEP06\(2020\)171](https://doi.org/10.1007/JHEP06(2020)171) arXiv:2003.05359 [hep-th]
- M.R. Garousi, H. Razaghian, Phys. Rev. D **100**(10), 106007 (2019). <https://doi.org/10.1103/PhysRevD.100.106007> arXiv:1905.10800 [hep-th]
- T.H. Buscher, Phys. Lett. B **194**, 59 (1987). [https://doi.org/10.1016/0370-2693\(87\)90769-6](https://doi.org/10.1016/0370-2693(87)90769-6)
- T.H. Buscher, Phys. Lett. B **201**, 466 (1988). [https://doi.org/10.1016/0370-2693\(88\)90602-8](https://doi.org/10.1016/0370-2693(88)90602-8)
- M.R. Garousi, Phys. Lett. B **809**, 135733 (2020). <https://doi.org/10.1016/j.physletb.2020.135733> arXiv:1907.09168 [hep-th]
- Y. Akou, M.R. Garousi, Eur. Phys. J. C **81**(3), 201 (2021). <https://doi.org/10.1140/epjc/s10052-021-08990-4> arXiv:2012.13264 [hep-th]
- R.C. Myers, Phys. Rev. D **36**, 392 (1987). <https://doi.org/10.1103/PhysRevD.36.392>
- T. Nutma, Comput. Phys. Commun. **185**, 1719 (2014). <https://doi.org/10.1016/j.cpc.2014.02.006> arXiv:1308.3493 [cs.SC]
- N. Kaloper, K.A. Meissner, Phys. Rev. D **56**, 7940 (1997). <https://doi.org/10.1103/PhysRevD.56.7940> arXiv:hep-th/9705193
- R. Borsato, A. Vilar López, L. Wulff, JHEP **07**(07), 103 (2020). [https://doi.org/10.1007/JHEP07\(2020\)103](https://doi.org/10.1007/JHEP07(2020)103) arXiv:2003.05867 [hep-th]
- O. Hohm, B. Zwiebach, JHEP **04**, 101 (2016). [https://doi.org/10.1007/JHEP04\(2016\)101](https://doi.org/10.1007/JHEP04(2016)101) arXiv:1510.00005 [hep-th]
- O. Hohm, B. Zwiebach, Phys. Rev. D **100**(12), 126011 (2019). <https://doi.org/10.1103/PhysRevD.100.126011> arXiv:1905.06963 [hep-th]